



POST HARVEST ROT OF FRUITS AND VEGETABLES

DISSERTATION SUBMITTED TO THE
ALIGARH MUSLIM UNIVERSITY ALIGARH
IN PARTIAL FULFILMENT OF THE REQUIREMENTS
FOR THE DEGREE OF

Master of Philosophy IN **BOTANY**

By
Pammie Joshi

DEPARTMENT OF BOTANY
ALIGARH MUSLIM UNIVERSITY
ALIGARH

1984



DS564

ACKNOWLEDGEMENT

I would like to convey my profound gratitude to Dr. S.K. Saxena, Chairman of the Department of Botany, Aligarh Muslim University, Aligarh, for his invaluable guidance and supervision. Without his cooperation this dissertation could have not taken this shape.

I am equally deeply thankful to Dr. Abrar M. Khan, Professor Emeritus for giving his valuable ideas from time to time. I am also thankful to Sri Mahendra Pal Sharma who took great pain to type this dissertation very neatly. At last but not the least, I must not forget to show my gratitude to my parents and sister who helped me in various ways.



(Pammie Joshi)

C O N T E N T S

INTRODUCTION AND REVIEW OF LITERATURE	1
FUTURE PLAN OF WORK	35
MATERIALS AND METHODS	38
REFERENCES	51

INTRODUCTION AND REVIEW OF LITERATURE

INTRODUCTION AND REVIEW OF LITERATURE

Fruits and vegetables form an important component of the diet of vegetarians providing sources of vitamins and minerals. This is all the more true for India where a majority of the population is vegetarian.

Although exact data on the annual production of fruits and vegetables in India are not available but it is estimated that acreage under cultivation ranges between 28-50 million acres. The produce is not sufficient enough for consumption of the people partly because major part is destroyed by pests and diseases including post harvest ^{wast} during handling, transportation and storage.

Powell (1906) named this discipline as Market pathology, dealing with diseases affecting fresh fruits and vegetables after the harvest and before their consumption. Broadly speaking the science is concerned with the spoilage occurring during the operations of harvesting, packing, handling, transportation, storing the produce.

According to Bourne (1976) the loss of food during post harvest can be defined as "that weight of wholesome edible product (exclusive of moisture content) that is normally consumed by human and that has been separated from the medium and the site of its immediate growth or production by deliberate human action with the intention of using it for human feeding but which for any reason fails to be consumed by humans".

Over all losses to fruits and vegetables from the farms to the consumers are about 12% (Columbic, 1964, Lentz & Varsden Berg 1971) of the total production. According to U.S.D.A. estimates (1965) losses of the kind in fruits and vegetables are to the tune of 23%.

In India Chemulu & Thakur (1968) while conducting a survey of the extent of losses estimated that potatoes suffer a loss of 24.0%, tomato 19.3%, mango 17.7%, banana 13.5% and apple 10.8%. Harvey (1978) while reviewing the literature of post harvest losses pointed out that the total percent loss in apples were 1.7-3.6, cucumber 2.9-7.9, grapes 5.0-10.3, lettuce 4.1-11.7, oranges 3.2-4.2, peaches 2.3-12.6, pears 4.1 - 10.1 and tomato 11.4 - 14.2. By and large the losses are much more in developing countries like India than in developed countries because of lack of advanced technology. Onosirosan and Fatunla (1976) showed that up to 21% of the potential harvest of tomato fruits were lost to rots in the field and an additional 5 - 20% rotted during marketing. Parpia (1976) on the other hand observed that losses during storage and handling of all crops in tropical Africa were to the extent of more than 30%. Thus it is now realised that post harvest diseases cause considerable losses to fruits and vegetables round the globe.

Large number of post harvest disease of various fruits and vegetables have been reported from time to time and some of the important ones are listed in Table I.

TABLE - I

List of some of the important post harvest diseases of
fruits and vegetables.

<u>Name of the fruit/ vegetable</u>	<u>Name of the fungus</u>	<u>References</u>
<u>1</u>	<u>2</u>	<u>3</u>
<u>Aegle marmelos</u> (Bel)	<u>Fusarium solani</u> (Mart) App & Wr	Sharma <u>et al</u> (1979)
	<u>Phoma glomerata</u> (Corda) Wr & Hochapfel	Pandey <u>et al</u> (1980)
<u>Annona squamosa</u> L.	<u>Glomerella cingulata</u>	Dhinra <u>et al</u> (1981 b)
<u>Averrhoa carambola</u> L. (Kamrakh)	<u>Cladosporium herbarum</u> (Pers) Link	Sharma & Khen (1978)
<u>Carica papaya</u> L.	<u>Pythium aphanidermatum</u>	Dhargava (1941)
	<u>Rhizopus stolonifer</u>	Tandon & Mishra (1969)
	<u>Fusarium solani</u> (Mart) App & Wr	Quimo & Quimo (1978)
	<u>Corynespora cassicola</u>	Quimo & Abilay (1981)
	<u>Phytophthora capsii</u>	Aragaki & Uchida (1979)
	<u>Colletotrichum capsici</u> (Syd) Butler & Bisby	Lai <u>et al</u> (1980)
	<u>Aspergillus niger</u> van Tiegh	Garcha & Singh (1980)
	<u>Trichothecium roseum</u> Link	Saxena & Jain (1981)
<u>Carissa</u> <u>Carandas</u>	<u>Myrothecium roridum</u> Tode ex Fr	Kumar & Tandon (1979)
	<u>Cochliobolus spicifer</u> Nelson	Wadia & Manoharachary (1979)
	<u>Rhizoctonia solani</u> Kuehn	
<u>Citrullus</u> <u>lanatus</u>	<u>Fusarium solani</u> (Mart) App & Wr	Ikediugwu & Ogileva (1979)
	<u>Corticium rolfsii</u>	Ikediugwu (1981)

1	2	3
<u>Citrullus</u> <u>vulgaris</u>	<u>Fusarium oxysporum</u> (Sohl ex Fr) em Snyder	Mathur & Mathur (1958) Suryanarayana & Nath (1963)
	<u>Fusarium equiseti</u> (Corda) Sacc	Cheurasia (1980)
<u>Citrus</u> sp	<u>Aspergillus niger</u> van Tiegh	Srivastava & Tandon (1969)
	<u>Trichothecium roseum</u> Link	Cheema & Jeyarajan (1972)
	<u>Geotrichum candidum</u> Link ex Pers	Kamal et al (1978)
	<u>Phytophthora citrophthora</u> <u>P. (nicotianae var) para-</u> <u>sitica</u> (Dastur) <u>P. syringae</u> <u>P. hibernalis</u>	} Feld et al (1979)
	<u>Trichoderma lignorum</u> (Tode) Harz	Kanaujia (1979)
	<u>Penicillium digitatum</u> Sacc <u>P. italicum</u> Welmeyer <u>Botrytis cinerea</u> (Pers) Link <u>Sclerotinia sclerotiorum</u> (Lib) de Bary	} Bai (1979)
	<u>Penicillium digitatum</u> Sacc <u>P. italicum</u> Welmeyer <u>Diplodia natalensis</u> (Bo- tryodiplodia theobromae) <u>Colletotrichum gloeospori-</u> <u>idea</u>	} Kumbhare and Choudhari (1980)

1	2	3
	<u>Alternaria alternata</u> (Fr) Kofasler	Garcha & Singh (1980)
	<u>Aspergillus fumigatus</u> Thom	
	<u>A. niger</u> van Tiegh	
	<u>Botryodiplodia theobromae</u> Pat	
	<u>Fusarium semitectum</u> Berk & Rav	
	<u>Penicillium crustosum</u>	
	<u>P. cyclosporum</u>	
	<u>Rhizopus stolonifer</u>	
	<u>Aspergillus aculeatus</u> Lis	Kanaujia (1981 a)
	<u>Trichoderma viride</u> Pers ex Fr	Kanaujia (1981 b)
	<u>Hormonema dematioides</u>	Rajak & Gautam (1981)
	<u>Drechslera</u> (Sclerosphaeria) <u>rustrata</u>	
	<u>Acremonium oycosum</u>	
	<u>Colletotrichum fraseriae</u> Brooks	Howard (1972)
<u>Fraseria</u> <u>virginiana</u> (Strawberry)	<u>Dendrophoma obscurans</u> (Ell & Ev)	Howard & Albregts (1973)
	<u>Pestalotia longisetula</u>	Howard & Albregts (1974a)
	<u>Alternaria tenuissima</u> (Nees ex Fr)	Howard & Albregts (1974b)
	<u>Phytophthora nicotianae</u> Var <u>parasitica</u> (Dastur)	Matsuzaki et al (1981) Suzvi et al (1981)
	<u>Phytophthora caetorum</u> (Leb & Cohn)	Kao & Lev (1981)
	<u>P. citrophthora</u>	

1	2	3
	<u>Mucor</u> sp	} Dennis (1981)
	<u>Rhizopus</u> sp	
	<u>Botrytis cinerea</u>	
<u>Litchi chinensis</u> (Litchi)	<u>Geotrichum candidum</u> Link ex. Fr	Jamaluddin <u>et al</u> (1978)
<u>Malus</u> (Apple)	<u>Trichothecium roseum</u> (Pers) Link ex Fr	Agarwala & Sharma (1968)
		Sree Kantilal <u>et al</u> (1974)
	<u>Uromyces</u> sp	Jones & Fields (1972)
	<u>Pestalotia</u> sp	Nathur & Agarwal (1974)
	<u>Glomerella cingulata</u>	Brook (1978)
	<u>Aspergillus candidus</u> Link	Thind <u>et al</u> (1978)
	<u>Phytophthora syringae</u>	Bolay (1978)
		Ustone (1979)
	<u>Monochaetia mali</u>	Naik (1979)
	<u>Clathridium corticola</u>	Thind <u>et al</u> (1979)
	<u>Penicillium expansum</u> Link	} Borecka (1979)
	<u>P. diversum</u>	
	<u>P. cyclopium</u> Westling	
	<u>P. spinulosum</u> Thom	
	<u>Corticium centrifugum</u>	Veresub & Ilman (1980)
	<u>Fusarium solani</u> (Mart) App & Wollenw	} Sumbali <u>et al</u> (1980)
	<u>F. equiseti</u> (Corda) Sacc	
	<u>Penicillium expansum</u> Link	} Babovic <u>et al</u> (1980)
	<u>Stemphylium botryosum</u> (Plo- ospora herbarum) Wallr	

1	2	3
	<u><i>Eusporium lactis</i></u>	
	<u><i>Monilia</i> (<i>Sclerotinia</i>)</u>	
	<u><i>fructigena</i></u>	
	<u><i>Gleosporium fructigenum</i></u>	
	(<i>Glomerella cingulata</i>)	
		Bobovic et al (1980)
	<u><i>Calcariosporium</i> sp</u>	
	<u><i>Rhizopus oryzae</i> Went &</u>	
	<u>Goerlinga</u>	
	<u><i>Trichothecium roseum</i> Link</u>	
		Garcha & Singh (1980)
	<u><i>Penicillium expansum</i> Link</u>	
	<u><i>Cladosporium herbarum</i> (Pers)</u>	
	<u>Link</u>	
		Eiche (1981)
	<u><i>Gleosporium</i> sp</u>	
	<u><i>Botrytis cinerea</i> Pers</u>	
	<u><i>Penicillium</i> sp</u>	
	<u><i>Nectria gallicola</i> Bers.</u>	
	<u>Bark</u>	
		John & Burth (1981)
	<u><i>Botryodiplodia theobromae</i></u>	
	<u>Pat</u>	
		Bhargava et al (1981)
<u><i>Mangifera indica</i></u>	<u><i>Phycolopora rhodina</i> (B & G)</u>	
	<u><i>Hendersonia creberrima</i></u>	
		Brodrick & Westhuisen (1977)
	<u><i>Phomopsis mangifera</i> Ahmad</u>	
		Amud-Laxminarayana & Reddy (1978)
	<u><i>Aspergillus niger</i> van</u>	
	<u>Tiegh</u>	
	<u><i>A. terreus</i></u>	
	<u><i>A. flavus</i></u>	
	<u><i>A. fumigatus</i> Fres</u>	
		Garcha & Singh (1980)
	<u><i>Botryodiplodia theobromae</i></u>	
	<u>Pat</u>	
	<u><i>Botryosphaeria ribis</i></u>	

1	2	3
	<u>A. flavus</u> Link	Najaudar & Modi (1981)
<u>Musa</u> (Banana)	<u>Thielaviopsis paradoxa</u> (De Seynes)	Garcia (1972)
	<u>Botryodiplodia</u> sp	Lantican & Quinio (1978)
	<u>Fusarium moniliforme</u> Sheldon	} Khanna & Chandra (1979)
	<u>F. roseum</u> Link	
	<u>Fusarium moniliforme</u> (Gibber- ella fujikuroi) Sheldon	Abdel-Sattar <u>et al</u> (1979)
	<u>Ceratocystis paradoxa</u> (de Seynes) Noreau	Abdel-Sattar <u>et al</u> (1979)
	<u>Cylindrocarpum</u> (Nectria) <u>radialis</u>	} Sharma & Khan (1981)
	<u>Aspergillus terreus</u> Thom	
<u>Phyllanthus</u> <u>emblica</u> L. (Aonla)	<u>Phoma emblicae</u> Jamaluddin	Jamaluddin <u>et al</u> (1979)
	<u>P. putaminum</u> Speg	Pandey <u>et al</u> (1980)
	<u>Cladosporium tenuissimum</u> Cooke	} Jamaluddin (1981)
	<u>C. cladosporioides</u> (Pres) de vries	
<u>Physalis</u> <u>peruviana</u>	<u>Fusarium equiseti</u>	Rao & Subramaniam (1979a)
Cape goose- berry or Raspberry	<u>Drechslera</u> (Setosphaeria) <u>rostrata</u>	} Tandon & Singh (1981b)
	<u>Alternaria alternata</u> (Fr) Keissler	
	<u>Curvularia lunata</u> (Cochlibolus lunatus) var-aeria	
<u>Prunus</u> <u>avium</u> (Cherry)	<u>Alternaria dendritia</u>	Gupta & Verma (1980)
	<u>Trichothecium</u> sp	} Bhargava <u>et al</u> (1981)
	<u>Rhizopus</u> sp	
	<u>Alternaria</u> sp	
	<u>Fusarium</u> sp	

1	2	3
<u>P. domestica</u> (Plum)	<u>Monilinia fructicola</u> (Wint) Honey	Kable (1973)
	<u>Sclerotinia fructicola</u>	Tate & Corbin (1979)
	<u>Glomerella cingulata</u>	Maiti <u>et al</u> (1981)
<u>P. persica</u> (Peach)	<u>Phytophthora syringae</u> (Kleb) Kleb	D' ercole & Flori (1978)
	<u>Sclerotinia fructicola</u>	Tate & Corbin (1979)
	<u>Monilia fructicola</u> (Wint)	Phillips & Harris (1980)
<u>Psidium guajava</u> (Guava)	<u>Macrophoma allahabadensis</u>	Kapoor & Tandon (1970)
	<u>Diplodia natalensis</u>	Rajagopalan & Wilson (1972)
	<u>Mucor heimalis</u> Welmeyer	Kunimoto <u>et al</u> (1977)
	<u>Phytophthora nicotianae</u> var Parasitica	Singh <u>et al</u> (1978)
	<u>Pestalotia olivacea</u> Guba	Dhingra & Melrotra (1980)
	<u>Aspergillus nidulans</u> var	Garcha & Singh (1980)
	<u>Rhizopus stolonifer</u> (Ehrhnb ex Fr)	Ooka (1981)
<u>Punica granatum</u> L. (Pomegranate)	<u>Aspergillus flavus</u> var	Srivastava & Tandon (1971)
	<u>Coniella novine - zelandiae</u>	} Sharma & Jain (1979)
	<u>C. granati</u>	
	<u>Aspergillus niger</u> van Tiegh	Philip (1981)
<u>Pyrus Communis</u> L. (Pear)	<u>Coremiella cubispora</u>	Lucas <u>et al</u> (1977)
	<u>Sclerotinia fructigena</u>	} Byrde & Willetts (1977)
	<u>S. fructicola</u>	
	<u>S. laxa</u>	
	<u>S. laxa</u> f. sp. mali	

1	2	
	<u>Sclerotium rolfsii</u> Sacc	Sunabali & Mehrotra (1980)
	<u>Trichoderma viride</u> Pers Ex Fr	Roy & Sharma (1980)
	<u>Penicillium crustosum</u>	Garcha & Singh (1980)
	<u>Phoma glomerata</u> (Corda) Wr & Hochapfel	Chohan & Chand (1981)
	<u>Penicillium expansum</u> Link	Borecka (1979)
	<u>P. diversum</u>	
	<u>P. cyclopium</u> Westling	
	<u>P. spinulosum</u> Thom	
<u>Vitis</u> <u>vinifera</u> L. (Grape)	<u>Botrytis cinerea</u> Pers	Maclellan & Hewitt (1973)
	<u>Greenaria fuliginosa</u> Scribner & Viala	Prakash et al (1976)
	<u>Guignardia bidwellii</u> (Ellis) Viala & Raven	Ferrin & Remadell (1978)
<u>Zizyphus</u> <u>auritiana</u> L. (Ber)	<u>Phoma hispanica</u>	Gupta & Madan (1979)
	<u>Alternaria</u> state of <u>Pleospora</u>	
	<u>Colletotrichum gloeospori-</u> <u>oides</u>	
	<u>Trichothecium roseum</u>	
	<u>Alternaria alternata</u> (Fr) Keissler	Wadia & Manoharachry (1981)
	<u>Fusarium decemcellulare</u> (Calonectria rigidiuscula)	
	<u>Cladosporium cladosporioides</u> (Fres) de vries	
<u>Allium</u> <u>cepa</u> (Onion)	<u>Sclerotium cepivorum</u> Berk	Rondomanski (1978)
	<u>Aspergillus niger</u> van Tiegh	Sharma & Roy
	<u>Cephalosporium</u> (Acremonium) <u>curtipes</u>	

1	2	3
	<u>Pleospora herbarum</u>	Wu (1978)
	<u>Sclerotinia sclerotiorum</u>	Teupkova (1979)
	<u>Aspergillus niger</u> van Tiegh	Raju & Raj (1981)
<u>Allium</u> <u>sativum</u> L. (Garlic)	<u>Fusarium solani</u> (Mart) App et Wt emend	Nicoll et al (1972)
	<u>Cephalosporium</u> (Acremonium) <u>curtipes</u>	Roy et al (1978)
	<u>Penicillium canum</u>	} Georgieva & Kotev (1979 b)
	<u>P. corymbiferum</u>	
	<u>Fusarium oxysporum</u>	} Wu (1979)
	<u>F. solani</u>	
	<u>Fusarium oxysporum</u> f. sp cavae	} Georgieva & Kotev (1979 c)
	<u>Penicillium canum</u>	
	<u>P. corymbiferum</u>	
	<u>Helminthosporium allii</u>	
<u>Amorphosa-</u> <u>llus campe-</u> <u>mulatus</u> Blume	<u>Sclerotium rolfsii</u> Sacc	Tyagi & Sharma (1978)
<u>Benincasa</u> <u>hispida</u> (Petha fruit)	<u>Fusarium solani</u>	} Sharma et al (1981)
	<u>F. moniliforme</u> (Gibberella fujikuroi)	
	<u>Penicillium citrinum</u>	Roy et al (1981)
<u>Capsicum</u> sp	<u>Colletotrichum capsici</u>	Singh et al (1978)
	<u>Alternaria alternata</u> (Fr) Keiseler	Shaukat et al (1979)
	<u>Fusarium solani</u>	} Nicosa & Ilag (1979)
	<u>F. oxysporum</u>	

1	2	3
	<u>Helminthosporium</u> (Broch-slera) <u>halodes</u>	} Laxminarayan & Reddy (1980)
	<u>Myrothecium</u> <u>roridum</u>	
	<u>Pestalotiopsis</u> <u>thana</u>	
	<u>Aspergillus</u> sp	Seenappa et al (1981)
<u>Cucumis</u> sp	<u>Pythium</u> <u>butleri</u>	} Masfy et al (1978)
	A sterile phycomycete	
	<u>Cladoporium</u> <u>herbarum</u>	} Georgieva & Kotev (1979 a)
	<u>Penicillium</u> <u>italicum</u>	
<u>Daucus</u> <u>carota</u>	<u>Alternaria</u> <u>radicina</u>	} Varkhrushcheva (1973)
	<u>Stemphylium</u> <u>botryosum</u>	
	<u>Chalaropsis</u> <u>thieleioides</u>	Schaffrath (1978)
	<u>Sclerotinia</u> <u>sclerotiorum</u>	Tsupkova (1979)
	<u>Phytophthora</u> sp	Stelfox & Henry (1979)
	<u>Stemphylium</u> <u>radicium</u> (<u>Alternaria</u> <u>radicina</u>)	Glaser (1979)
	<u>Geotrichum</u> <u>candidum</u>	Suhag & Duhan (1981)
<u>Lycopersi-</u> <u>um escul-</u> <u>entum</u> (Tomato)	<u>Pythium</u> <u>ultimum</u>	} Pearson & Hall (1974) Sonoda (1974)
	<u>P. aphanidermatum</u>	
	<u>Geotrichum</u> <u>candidum</u>	} Barts (1981) Onesirosoan & Fatunla (1977)
	<u>Rhizoctonia</u> <u>solani</u>	
	<u>Sclerotium</u> <u>rolfsii</u>	
	<u>Pythium</u> <u>aphanidermatum</u>	
	<u>Botryodiplodia</u> <u>theobromae</u>	
	<u>Eusporium</u> <u>equiseti</u>	
	<u>E. xylarioides</u>	

1	2	3
	<u>Syncephalastrum racemosum</u>	Rao & Subramanian (1979b)
	<u>Fusarium solani</u> (Mart)	Sacc Garg & Gupta (1979)
	<u>Alternaria alternata</u> (Fr)	Tani <u>et al</u> (1979)
	Keissler	
	<u>Phoma destructiva</u>	} Kogan (1979)
	<u>Ascochyta</u> (Didymella)	
	<u>lycopersici</u>	
	<u>Microspora oxysae</u> (Berk	} Chary <u>et al</u> (1980)
	& Br) Petch	
	<u>Stomphylus vesicarius</u>	
	(Wallroth) Sacc	
	<u>Phytophthora nicotianae</u>	D' ercole (1981)
	var <u>nicotianae</u>	
	<u>Myrothecium carmichaeli</u>	Tandon & Singh(1981 a)
<u>Nomordica</u> <u>Charatia</u> (Bitter gourd)	<u>Alternaria tenuissima</u>	} Sharma & Bhargava (1979)
	<u>Myrothecium roridum</u>	
	<u>Glomerella cingulata</u>	Verma <u>et al</u> (1981)
<u>Solanum</u> <u>tuberosum</u>	<u>Phytophthora erythrae-</u>	Carroll & Sasser (1974)
	<u>ptica</u> Pethybr	Lopkova (1976)
	<u>Ophiobolus porphyrogonus</u>	Popkova & Kovalova (1974)
	(Tode) Sacc	
	<u>Acremonium atra</u> (Corda)	El-Khadem & El-Kassas
	Sacc	(1978)
	<u>Fusarium solani</u> (Mart)	Sacc Braslavsky (1978)
	<u>Aspergillus flavus</u> Link	Gupta <u>et al</u> (1978)
	<u>Fusarium caeruleum</u> (Lib)	Langerfeld (1979)
	Sacc	
	<u>Phoma solanicola</u> Prill &	Popov (1979)
	Dell	
	<u>Fusarium trichothecioides</u>	Marte & Tamburi (1980)
	Woll	

1	2	3
	<u>Fusarium equiseti</u> (Corda) Sacc	Rai (1980)
	<u>Phoma cupyrena</u> Sacc	Langerfeld (1980)
	<u>Phoma</u> spp	Janke & Zott (1981)
	<u>Fusarium</u> sp	Gusev (1981)
<u>Solenum melongena</u> L.	<u>Phomopsis vexans</u> (Sacc et Syed) Darter	Panwar et al (1972) Chowdhury & Hasiza (1981)
	<u>Chrysosporium oryzae</u> (Gilman & Abbott)	Dhingra & Mehrotra (1980)
	<u>Rhizopus nodosus</u>	Vyas et al (1980)
	<u>Fusarium moniliiforme</u> Sheld Darter	(1981)
	<u>Rhizopus oryzae</u> Went & Geerlings	Ali & Shukla (1981)
<u>Trichosanthis ananassa</u> (Snake Gourd)	<u>Penicillium janthinellum</u>	Rao & Thirupathaiah (1980)
<u>I. dioica</u> Roxb (Pointed gourd)	<u>Fusarium equiseti</u> (Corda) Sacc	Kritagyan & Singh (1980)
	<u>Alternaria alternata</u> (Fr) Keissler	
	<u>Glomerella singulata</u> (Strom) Spauld & Schrenk	
	<u>Helminthosporium</u> sp	
	<u>Curvularia lunata</u> (Walker)	
	<u>Doodya var acris</u> (Batista) Lima, & Vasconcelos) Ellis	
<u>Zingiber officinale</u> (Ginger)	<u>Diplodia natalensis</u>	Wilson & Balagopal (1973)
	<u>Pythium deliense</u>	Haware & Joshi (1976)

BIOCHEMICAL CHANGES:-

Fungal invasion brings about changes in chemical constituents of the fruits. There is a great shift in the concentration of carbohydrates, organic acids, phenols, aminoacids, vitamins etc. (Ton'ze 1964, Srivastava & Tandon 1966, Stretch & Cappellini 1965, Aulakh et al 1970a, 1970b).

Carbohydrates:- Drastic changes in sugar content has been observed in apples infected with Aspergillus terreus Thom (Chandra & Tandon, 1963), guava, sapodilla, papaya and banana fruits with Pestalotia psidii P. Henn, Colletotrichum papayae P. Henn, Fusarium sp, Phoma psidii, Gleosporium sp, Botryodiplodia theobromae Pat, and Pestalotia sapodilla P. Henn (Ghosh et al 1964) cucumber by Pythium aphanidermatum (Eda) Fitz (Mc Comb & Winstead, 1964).

Ghosh & Tandon (1965a) reported the presence of oligosaccharides in mango fruits infected with Colletotrichum gloeosporioides Pens; in musambi infected with Botryodiplodia theobromae Pat. Srivastava & Tandon (1966) reported that oligosaccharides developed on the 4th day after infection. Oligosaccharides were also detected as a result of infection with Pestalotia sapota P. Henn & B. theobromae Pat in sapota & with Gleosporium psidii Delacr (L. Fr) on a variety of guava (safeda). However, infection of Pestalotia psidii P. Henn & Phoma psidii on guava did not form oligosaccharides (Tandon, 1967).

Banana infected with Botryodiplodia theobromae Pat showed loss in fructose and glucose content (Williamson & Tandon, 1965). Kapoor & Tandon (1967) reported decrease in glucose, fructose and sucrose content in guava fruits infected with Macrophomina alishahabensis Kapoor & Tandon, and musambi fruits infected with Aspergillus niger van Tiegh (Singh, 1968). Similar results have been observed in a tomato variety infected with Dreschlera australiensis (Bugn) Subram & Jain (Kapoor, 1968); papaya and banana fruits infected with Rhizopus stolonifer (Fr) Lind, tomato fruits infected with Phoma destructiva (Aulakh et al, 1970b).

Chahal & Grover (1972) while carrying out changes in the sugar content in Chilli fruits infected with Choanophora cucurbitarum (Berx & Rav) Thaxter, found that in the healthy fruits two sugars viz. fructose, D-xylose were detected but after 3 days of pathogenesis both the sugars disappeared. Changes in sugar content have also been reported in banana fruits infected with Alternaria alternata (Fr) Keissler & Cochliobolus sp. epicifer Nelson (Prasad, 1974). tomato fruits infected with Cylindrocarpum scoparium Morgan, Colletotrichum sp Myrothecium toridum Tode ex Pers, Chilli fruits infected with Rhizopus stolonifer (Fr) Lind (Tandon et al, 1974), that in papaya infected with Phomopsis caricae papayae Petrax & Cif (Dhingra & Khare, 1975); tomato infected by Alternaria solani (Ellis & Mart) Jones & Grout & A. tenuis Nees ex Pers (Mehta et al, 1975) and musambi fruits infected with Botryodiplodia theobromae Pat (Ali, 1976). Singh & Chohan (1981) reported that the concentra-

-tion of 3-4 sugars which were detected in healthy fruits of cucurbits decreased when infected with Macrophomina phaseolina.

Organic acids- Organic acids like fumaric, tartaric, malic, succinic, oxalic acids which are the chief constituents of several fruits and vegetables also undergo changes as a result of infection with pathogens. Kapoor & Tandon (1969), Singh & Tandon (1970) Tandon, Srivastava & Jamaluddin (1974) reported that organic contents of fruits decreased in infections with various fungi.

Fumaric, malonic and citric acids which were observed in healthy fruits of tomato, were also observed in fruits infected with Drechslera australiensis (Dugn) Subram & Jain, but the content was very low except that of fumaric acid which showed an increase in the content. The production of succinic acid during pathogenesis of fruits is interesting (Kapoor & Tandon 1969). Tandon et al (1974) reported that chilli fruits infected with Rhizopus stolonifer (Fr) Lind, & tomato fruits infected with Colletotrichum sp, Cylindrocarpum acoperium Morgan, Myrothecium veridum Tode ex Pers showed a decrease in the organic acid content. Mehta et al (1977) carried out analysis of extracts of healthy and diseased tomato fruits infected with Alternaria solani (Ellis & Mart) Jones & Grout & A. tenuis Nees ex Pers & observed that there was considerable changes in organic acid content due to

pathogenesis. These observations led them to conclude that succinic acid was synthesized in the diseased fruits. Jamaluddin (1979) while studying changes in the organic acid content of healthy and diseased fruits of peach (Prunus persica) found that in the fruits infected with Aspergillus flavus the quantity of succinic, malonic and citric acids increased while the quantity of malic and tartaric acid decreased. Fumaric acid and an unknown acid with (RF 0.09) were detected only in the diseased fruits. In general the total amount of organic acids increased under pathogenesis.

Recently Majumdar & Modi (1981) while carrying out biochemical studies on spoilage of mangoes by Aspergillus flavus Link observed a high level of citric acid in infected fruits. Thind et al (1979) reported that the concentration of the 10 organic acids which were found in high quantity in healthy fruits showed a slight decrease in quantity after infection with Clathridium corticola in apple fruits.

Ascorbic Acids- It has been observed that the ascorbic acid content decreases when fruits are infected with fungal pathogens (Bhutani 1946).

Mango fruits infected with Colletotrichum gloeosporioides Penz & Aspergillus niger van Tiegh, and guava fruits infected with Pestalotia psidii P. Henn, Phoma psidii P. Henn & Gleosporium psidii P. Henn, (Ghosh et al 1965) showed decrease in ascorbic acid content. The rate of decrease of ascorbic acid was faster in mango fruits

infected with Botryodiplodia theobromae Pat (Srivastava & Tandon, 1966b). In musambi fruits which were infected with B. theobromae Pat no ascorbic acid was detected (Srivastava & Tandon 1966c). Chahal & Grover (1972) carried out studies on chilli fruits infected with Chenopodia cucurbitarum (Berk & Rav) Thaxter; and reported that ascorbic acid content of diseased chilli fruits declined rapidly after 5 days of pathogenesis.

Significant losses in ascorbic acid content was also observed in papaya fruits damaged by 6 Post harvest pathogens, namely Alternaria tenuis (A. alternata) Nees ex Pers, Chaetomium globosum Kunze & Sohn, Curvularia lunata (Cochliobolus lunatus) (Walker) Boedij, Cylindrocarpum tonkinense, Fusarium oxysporium (Schl ex Fr) Syd et Hans & Helminthosporium spiciferum (Cochliobolus spicifer) (Prasad & Verma, 1978). Agrawal & Bisen (1978) reported that when Aspergillus niger van Tiegh & Alternaria tenuis (A. alternata) Nees ex Pers isolated from apple were inoculated into 5 apple cultivars, decrease in ascorbic acid was seen and this was much faster the apples inoculated with Aspergillus niger van Tiegh than Alternaria alternata Nees ex Pers. Fruits of litchi infected with Fusarium semitectum Berk & Rav, Alternaria alternata (Fr) Reissled & Cylindrocarpum tonkinense reduced the ascorbic acid content by 87.4 - 90.7% although some loss was also observed in control fruits which may be due to over ripening (Prasad & Sinha 1979). Decrease in ascorbic acid content during pathogenesis of lemon fruits with Fusarium oxysporum was

reported by Nararia & Reddy (1980).

Phenols:- Phenolic compounds are also one of the important constituents of a plant cell. These compounds provide resistance in fruits and vegetables. Higher concentration of phenolics in a variety of tomato resistant to Alternaria solani (Ellis & Mart) Jones & Grout was reported by Bhatia et al (1972). Similarly in tomato fruits infected with Alternaria tenuis Nees ex Pers & A. solani (Ellis & Mart) Jones & Grout were found to contain higher amount of phenols, (Mehta et al 1975).

A phenolic substance (RF value .32) was detected in the healthy fruits which decreased during pathogenesis by Alternaria solani and was totally lost in 12 days old infections. Two phenols (RF. .3 & .15) were observed in healthy host tissue of apple fruits but after infection with Clathridium corticola, the phenol (RF. .3) increased (Thind et al 1979). Post infection biochemical changes in apple fruits rotted by Penicillium expansum Link, Glomerella cingulata Spauld & Schen Trichothecium roseum Link, Monilinia (Sclerotinia) laxa & Rhizopus stolonifer (Fr) indicated a reduction in the phenolic content (Kaul & Munjal 1981).

Amino acids:- Post infection changes in amino acids have been studied in seasonal fruits such as in melons infected with Colletotrichum lagenarium (Pass) Ell & Ever (Tonze, 1964); Papaya with Gleosporium papayae P. Henn, Botryodiplodia

theobromae Pat, Colletotrichum pasayae P. Henn, Rhizopus nigricans Ehrenb; guava fruits with B. theobromae Pat, Phoma psidii P. Henn, Sclerotium psidii P. Henn, Pestalotia psidii P. Henn (Tandon, 1967); banana fruits with B. theobromae Pat, Sclerotium musarum Cooke & Mass.

Mango, sapota and citrus fruits infected with B. theobromae Pat, tomato with Phoma destructiva Plowder (Anilakh et al 1970a) showed significant changes in the amino acid content. Chahal & Grover (1972) carried out studies on healthy chilli fruits and also fruits infected with Chasmodon cucurbitarum (Berk & Rav) Thaxter. They detected eight amino acids namely Aspartic acid, glutamic acid, L-Alanine, D-Alanine, tryptophane, Arginine, histidine, glycine in healthy fruits but during pathogenesis all except glycine were reduced considerably. Tomato fruits infected with Cylindrocarpus scoparium Morgan, Myrothecium roridum Tode ex Pers, Colletotrichum sp, Alternaria tenuis Nees ex Pers, A. solani (Ellis & Mart) Jones & Grout (Mehta et al 1975 a,b) also exhibited a sharp decline in the amino acid content during pathogenesis.

Mehta et al (1977) carried out analysis of extracts of healthy and diseased tomato fruits infected with Alternaria solani (Ellis & Mart) Jones & Grout & A. tenuis Nees ex Pers & observed changes in the composition of aminoacids. The content of free aminoacids was considerably altered qualitatively and five aminoacids appeared in the diseased fruits. Singh & Chohan (1979) carried out experi-

-ments on changes in free aminoacids in fruits of bottlegourd, muskmelon & watermelon due to infection of Pythium butleri. Thind et al (1979) observed post infection changes in amino acid contents in apple fruits rotted with Clathridium corticola. Of the 10 free amino acids detected in the healthy tissues of apple fruits histidine, arginine, aspartic acid, alanine, leucine, valine, phenylalanine were completely utilized or disintegrated by the pathogen during pathogenesis. However, three newly synthesized aminoacids viz. Cystine, aminobutyric acid and tyrosine were detected in the infected tissues. Loss in amino acid content was also observed in tomato fruits infected by Phoma destructiva Plover (Hasija & Batra, 1981).

Kaul & Munjal (1981) determined biochemical changes including changes in aminoacids in apple fruits inoculated with Penicillium expansum Link, Monilia (Sclerotinia) laxa & Rhizopus stolonifer (Fr) Lind and obtained very interesting results.

Factors affecting post harvest losses- Large number of factors have been found to influence these losses in fruits and vegetables during storage and transits. Broadly they are classified as physical, chemical and biological factors.

Physical- This category includes (a) the initial quality of the commodity (b) the temperature at which the product is

held during handling, storage, transport and distribution (c) the h. . of the post harvest environment (d) the use of controlled or modified atmospheres during storage and transit (e) chemical treatments for the control of decay (f) heat treatments for decay control (g) packing and handling systems.

Initial quality- Fruit harvested at an immature stage may never ripen properly and may consequently constitute a post harvest loss when rejected at the retail or consumer level. Similarly fruits harvested at an over-mature stage may not survive the handling to which it is subjected during marketing. Thus these fruits become a loss in the marketing sequence (Harvey 1978).

Harvey (1978) further pointed out that the initial quality of fruits should be free from infections to avoid further rotting during storage and transport. This post harvest loss can be reduced by adopting an integrated approach of control methods (Harvey, 1978).

Temperatures- Normally for keeping the fruits and vegetables fresh, the storage is done at the temperature - in cold storage and in refrigerators. Parpia (1976) pointed out that lack of refrigeration capability is largely responsible for major post harvest losses in many developing countries. On the basis of experimentation, compilations have been made to indicate the optimum temperatures for maintenance of quality and prevention of losses during storage and transit.

(Asby, 1970, Ashare, 1974, Lutz & Gordenbury 1968, Pantastico 1974, Redit 1969, Whiteman 1957). MacLeod et al (1976) pointed out that there is a relationship between post harvest losses and temperature. These authors reported that lettuce become unsalable in only 12 days at 15°C but not until 35 days at 0°C. Asparagus was found more perishable as it became unsalable in only 5 days at 15°C but not until 35 days at 2°C. Storage of asparagus at 0°C is not permissible as it brings about chilling injury.

Lutz & Gordenbury (1968) showed that many tropical fruits such as mangoes and bananas are damaged if stored at a temperature below 7.5°C. Similarly green vegetables are also susceptible to freeze injury and are, therefore, recovered for storage at temperature above 7.5°C.

Temperature during storage has another problem of control of microorganisms. Certain microorganisms thrive best and are capable of growing even at low temperatures.

Studies on 11 species of Rhizopus, which is a very important pathogen of fruits and vegetables, were carried out by Harter & Weimer (1922) to determine the optimum temperature required by the species to grow. Rhizopus autogari Kocib, R. nigricans Ehrenb, grow best at 20-25°C. R. Oryzae Fischer & R. nodus Nannysolwski required 30°C as optimum temperature. R. Chiensis Ehrenb developed at 35°C.

Karmakar & Joshi (1940), Sethi (1948) and Mukerjee (1961) reported that below 10°C pathogens failed to grow on mango. A temperature range of 20 - 25°C was found to be the optimum temperature for the growth of Alternaria tenuis Nees ex Pers on Pears (Tandon & Ghosh 1962).

Tandon (1967) reported that Gleosporium musarum Cke & Mass caused least decay on banana fruits when stored at 10°C. Pestalotia sapotae P. Henn & Botryodiplodia theobromae Pat could not cause any decay on sapodilla at 10 - 15°C but considerable losses were observed at temperature between 25 - 30°C (Tandon & Mishra, 1969). Similar studies to determine the optimum temperature for pathogens to grow were made on Litchi fruits infected with Aspergillus flavus Link, A. niger van Tiegh, A. quadricolus Thom & Raper, A. variegatus Thom & Raper, A. nidulans (Eidam) Wint, Cylindrocarpum tankinense Hui, Pestalotia sp & Colletotrichum gloeosporioides Pers which caused optimum rot at temperatures between 15 - 25°C.

Mehra et al (1977) while carrying out pathological studies on fruit rot of tomato caused by Alternaria solani (Ellis & Mart) Jones & Grout & A. tenuis Nees ex Pers reported that the maximum rotting occurred at 28°C. Alternaria sp., causing a fruit rot of Juglans regia failed to germinate at 45°C, with 30°C as the optimum temperature for germination (Verma et al 1977). Colletotrichum capsici (Syd) But & Bis causing fruit rot of chillies grew best at 30°C (Singh et al 1978). Potato

decay by Fusarium solani (Mart) App & Wali F. roseum Link & F. oxysporum (Schl ex Fr) Syd et Hans was minimum when tubers were stored at 19°C (Lee 1978). Sharma & Khan (1979) reported that optimum temperatures for the growth of Alternaria alternata (Fr) Keissler & Cladosporium cladosporioides on cape gooseberry were 20 & 25°C respectively. Fusarium solani (Mart) App & Wali & F. oxysporum (Schl ex Fr) Syd et Hans causing fruit rot of chillies grew and sporulated at 20 - 30°C. (Micoesa & Ilag, 1979). Blue mold on garlic in Bulgaria caused by Penicillium canum & P. corymbiferum developed best at 20 - 25°C and 25 - 30°C respectively (Georgieva & Kotev, 1979). Low temperature (5, 10, 15°C) were unfavourable for the initiation of fruit rot in lime fruits caused by Aspergillus niger Van Teigh while at 25, 30, 35°C rapid rot occurred within a few days (Khanna & Chandra 1979 a). Fusarium moniliformae (Gibberella fujikuroi) Sheldon & F. roseum Link causing banana fruit rot was maximum at 25 - 35°C (Khanna & Chandra 1979 b). Rot of cucumber fruits during transportation caused by Cladosporium herbarium (Pers) Link & Penicillium italicum developed at low temperature 8 - 10°C maintained in refrigeration cars (Georgieva & Kotev 1979).

Cylindrocylindrium clavatum causing potato tuber rot was less at 15°C and 20°C, with no measurable disease occurring at 10°C. (Bolkan et al, 1981). Chargava et al (1981) carried out studies on Botrydiploia rot of apple and reported maximum rot at 36°C and minimum at 20°C while no rotting

was observed at 10°C. Pathological studies regarding optimum temperature for the development of Phomopsis vexans (Sacc & Syd) causing rot of brinjal fruit carried out by Choudhary & Pasija (1981) indicated that 25°C was the optimum temperature for rotting.

Relative Humidity:- Like temperature relative humidity equally plays an important role in the development of post harvest diseases. It not only affects the moisture loss from fruits or vegetables as evidenced by shriveling or wilting, but also affects the activity of the decay causing organisms. Moisture loss is serious in commodities that are stored for a long time e.g. apples etc. or in those that may be stored only for short periods.

Fungi in general required high moisture level for their growth and metabolism. Generally 95 to 100 percent relative humidity supports highest growth of most of the fungi. Relative humidity below 80 to 85 percent are by and large inhibitory for them. Conidia of Monilinia fructicola (Wint) Honey and survive till a very long period at 15 percent RH (Naqvi and Good 1957).

Very high relative humidity generally favour the growth of organisms causing decay, but do not necessarily result in increased decay losses in fruits and vegetables (Harvey, 1978). In potatoes for example, high RH promotes curing and healing by suberization which reduces invasion by micro organisms (Arteschwager, 1927).

Recent studies on the effect of RH during storage of Brussels sprouts, cabbage, leeks showed that at low temperature there is less decay at 98 - 100 percent relative humidity than at 90 - 95 percent or lower. (Lents & van den Berg 1971, Pendergrass & Isenberg 1974). The rot of banana, apple, mango, tomato and potato by Rhizopus spp. was not very much influenced by RH. However, a RH of 50 percent or above allowed the rapid development of soft rot with moderate to heavy intensity (Thakur 1972).

Mehta et al (1977) reported 100 percent as the optimum relative humidity for the growth of Alternaria solani (Ellis & Hart) Jones & Grent & A. tenuis Nees ex Pers on tomato fruits. Khanna & Chandra (1979) reported that fruits of lime when stored at 90 - 100 percent RH showed 100 percent rot in nine days due to infection caused by Aspergillus niger Van Tiegh. Maximum rotting of banana due to Fusarium moniliforme (Gibberella fukuroi) Sheldon & F. roseum Link (Tandon & Chandra 1979) occurred when the relative humidity was 50 percent. High intensity of fruit rot of Pyrus communis due to Trichoderma viride occurred at 60 percent RH (Roy & Sharma, 1980). Potato tubers stored in polyethylene boxes were badly damaged by Fusarium sulphureum (Gibberella cyanogena) at higher RH (98 - 100 percent compared with 80 percent) (Langerfield, 1981). Maximum rotting of brinjal fruits due to Phomopsis vexans was seen at 55 percent RH (Choudhary & Hasija, 1981).

High relative humidity, when desired in storage is generally maintained by the use of a refrigeration system. Steam injectors or atomizers have also been used for increasing RH in storage (Gentry & Guillon, 1966). Packing of fruits with protective films creates an environment with high RH within the package and thus makes the fruit liable to more damage than in storage rooms (Hardenburg, 1971, 1973).

Controlled or modified atmospheres:- Controlled or modified atmospheres are equally essential for good storage of fruits. These conditions are to be supplemented with refrigeration to have a better effect (Dewey 1977). Dewey et al (1969) Manis et al (1971) studied the effect of controlled atmospheres for the storage and transport of perishable agricultural commodities. By and large, the controlled and modified atmospheres inhibit the activity of decay organisms. Griverson (1969) reported that fruits which undergo ripening after harvest are more sensitive to controlled atmospheres. Therefore, controlled atmospheres are effective for those fruits that cannot be stored at the optimum temperature.

Lettuce was found to be stored better at 0°C and is therefore usually transported at temperature of 4°C to 5°C. Reduction of the amount of O₂ in a rail, car or truck from a normal 21 percent to 2 to 5 percent slows down the respiration rate (Linton, 1967 and Morris & Lader, 1977).

Strawberries shipped by air are much above their optimum holding temperature of 0°C for the major part of the transit period (Harvey et al, 1966). Therefore maintenance

There was a reduction in decay of peach rots due to Monilia sp. Potrytis sp and Rhizopus sp by post harvest treatment with thiabendazole carbendazim, imazalil, iprodione, captafol & vinclozolin (Nombela et al 1980). White rot disease of garlic caused by Sclerotium cepivorum Berk was reduced when fungicides like vinclozolin, iprodione, benomyl & dicloran was applied as a bulb dust (Tamiotti 1981).

Heat treatments:- Heat treatment has been widely recommended because it controls organisms that have already penetrated the fruit as well as those located on the surface of the fruits. In a way this treatment is comparable to pasteurization of dairy products in that the temperatures and exposure times do not cause appreciable change in the texture, flavour or other characteristics of the product, but do inactivate the pathogens. Studies carried out by several workers have shown that care should be taken to control the temperature because high temperature may cause injury to the fruit. Papaya and peaches have been treated with heat treatments to control fruitfly and pathogens such as Colletotrichum in Hawaii. Treating peaches for 3 - 5 minutes in water at 49°C or 1.5 minutes at 54°C controlled both brown rot (Monilia fructicola) and Rhizopus rot (Rhizopus stolonifer) (Smith et al 1964). A 2 to 3 minutes exposure to 52°C in water was most effective for control of fruits decay, without causing injury (Smith & Kedit, 1968). Akamine and Arisumi (1953) and Akamine and Goo (1969) suggested a treatment of

papaya for 20 minutes at 49°C for controlling anthracnose and other diseases. Akamine (1977) also suggested a 20 min dip in hot water at 47°C for controlling decay of mangoes. Wells (1971) recommended 1.5 min dips in water at 52°C for reducing the decay of nectarines by 65 to 75% compared to decay of untreated fruits. However, when hot water treatment was supplemented with treatment with Captan, benomyl, the effectiveness increased. This combination has another advantage that a very low concentration of the chemical was required as that when used alone. Spalding & Reader (1980) showed that post harvest decay was controlled by dipping mango fruits in hot water (52°C) containing .1% benomyl for 1-3 min.

Packing & Handling Systems:- Improper packaging and handling systems also aggravate the losses due to post harvest disease. Proper packaging of a product prevents not only bruising and crushing but can also reduce the moisture loss and pilferage in addition to maintaining a desirable environment during marketing.

According to Perkins (1959) packing of apples in polyethylene bags can reduce bruising caused during transport to the extent of 50%. Bags should be placed on their sides in layers within a partitioned master container rather than placing them upright through the full depth of the container.

Tight full packing of pears, plums, apricots, cherries and other tree fruits had resulted in reduction of losses from bruising. This kind of packing, however, would be different for different kinds of fruits and separate studies are needed for each type of fruits and vegetables. Fruits like cherries, peaches which are soft skinned may require some extra padding in packing (Gentry et al., 1965, Mitchell et al. 1964, 1965, 1967).

According to Guillev (1960), Olson et al. (1960), Patchen (1969), package design must be coordinated with cooling systems and refrigeration practices with in storage and in the transit taking into consideration the strength of the packing.

Box liners for cherries have been found effective in reducing the blue mold rot caused by Penicillium expansum LK ex Thom. McIntosh and Perritt (1958), proved that box liners caused an increase in decay because of the abundant moisture with in the packed boxes e.g. Bull's eye rot caused by Neofabraea pruniana Kieh was much higher in fruit stored in box liners than in those stored without liners. Hardenburg (1974) showed that treatment of fruits with certain fungicides before packing considerably reduced losses from decay. Box liners used for packing citrus when treated with biphenyl reduced the damage of citrus fruits from green mold rot caused by Penicillium digitatum, Sacc. and blue mold

rot caused by *P. italicum* Welmer (Rygg et al. 1964). According to Gerrhault (1955) the storage life of pears can be extended by 6 to 8 weeks with reduction in moisture loss by packing the fruit in sealed polyethylene liners. Singh & Gupta (1980) proved that decay of citrus fruits could be reduced when the fruits are stored in gunny bags or wooden boxes rather than, when stored in bamboo baskets.

Both grocers and consumers are responsible for reducing the post-harvest losses in perishable commodities. In addition to this it also involves each and every individual who handles the product during marketing.

It is, therefore, necessary to know the factors which influence the post harvest losses in fruits while they are stored in the consumers' place. Practically no work has been done on this aspect, more particularly in India.

- 35 -

FUTURE PLAN OF WORK

FUTURE PLAN OF WORK

It is clear from the brief review presented here that despite large number of studies there are lacunae in our knowledge which need to be filled in. Moreover nothing precisely is known what precautions to be taken to minimise the losses due to diseases in transit. Therefore, with this aim in view it is proposed to study the following:-

1. To isolate fungal pathogens from diseased fruits and vegetables viz. mango, apple, peach, banana, citrus, tomato, brinjal, potato, cressicum, carrot etc. and to diagnose them upto specific level.
2. To test their pathogenicity in fruits in the laboratory.
3. To study the effect of different types of semi solid and liquid media, different carbon and nitrogen sources and mineral elements on the growth of some of the known pathogenic fungi.
4. To study the effect of temperature, relative humidity and P^H on the development of pathogens isolated in 1 above.
5. To study the post-infection changes in the phenolic, aminoacid, carbohydrates, organic acid and ascorbic acid contents in diseased as well as healthy fruits and vegetables.
6. Effect of various substances on the production of enzymes.

7. To determine the effect of different packing systems on the development of post harvest losses.
8. To devise control measures.

MATERIALS AND METHODS

MATERIALS AND METHODS

Survey & Collection of diseased material:-

Diseased fruits and vegetables will be collected from the market and will be brought to the laboratory for isolating the fungi associated with the rot and identifying them.

Isolation, purification & maintenance of cultures:-

The diseased fruits and vegetables which will be collected from the local market will be kept in a moist chamber at $20 - 28^{\circ}\text{C}$, to facilitate the growth and sporulation of fungi.

For isolation of pathogens, the diseased fruits will be thoroughly washed with tap water and then the surface will be sterilized with .1% HgCl_2 solution by keeping them into the solution for 1 to 2 minutes. After repeated washing with distilled H_2O , diseased lesions will be cut from the junction of diseased tissues with the help of sharp sterilized knife. The diseased tissue will be transferred in sterilized petridishes containing PDA. Before transference of the diseased tissue the petridishes will be incubated at $28 \pm 1^{\circ}\text{C}$. Then the fungal hyphae growing out of the diseased tissues will be picked and transferred to fresh petri dishes containing PDA. The purified isolates will be maintained on PDA slants at lower temperatures. The isolates will be identified with the help

of available literature.

Pathogenicity test:-

Pathogenicity will be performed to confirm the pathogenic ability of the isolated pathogens. Uninjured, healthy specimens will be surface sterilized with .1% HgCl_2 solution and will be thoroughly washed with distilled water. The specimen will then be inoculated either by pricking, or by making a hole with the help of a sterilized cork borer (Granger and Horne, 1924). In both the cases the inoculum consisting of a mycelial agar disc will be cut from the margin of the freshly grown colony and will be placed over the incision made earlier. The inoculated specimen will be kept in moist chambers at $28 \pm 1^\circ\text{C}$. Uninoculated fruits will also be kept in the same conditions which will serve as control. Fruits will be observed daily and the symptoms will be compared with the original disease symptoms. Again the pathogens will be isolated and will be compared with the original cultures. Isolation and inoculation will be repeated many times so as to confirm Koch's postulates.

Effect of different Culture media:-

After identifying the fungi, the pathogen shall be cultured on different media, to find out the most suitable basal medium for further studies.

a) Potato - dextrose - agar medium:-

Peeled potatoes	200 g
Dextrose	20 g
Agar	20 g
Distilled Water	1000 ml.

b) Martin's Rose Bengal Streptomycin agar medium:-

Dextrose	10 g
Peptone	5 g
Potassium dihydrogen phosphate	..				1 g
Magnesium sulphate5 g
Rose Bengal (1 part in 30,000 parts of the medium)					
Agar	20 g
Streptomycin03 g
Distilled water	1000 ml.

c) Corn - meal - agar medium:-

Corn meal	20 g
Peptone (if desired)		20 g
Dextrose (if desired)		20 g
Agar	15 g
Distilled water	1000 ml.

d) Oat - meal - agar medium:-

Oats	100 g
Agar	15 g
Water	1000 ml.

e) Sabouraud's medium:-

Glucose (or maltose)		40 g
Peptone	10 g
Agar	15 g
Distilled water	1000 ml.

c) Czapek's medium-

Agar	20 g
Sucrose	30 g
Sodium nitrate	2 g
Potassium dihydrogen phosphate	..					1 g
Magnesium sulphate 7H ₂ O5 g
Potassium chloride5 g
Ferrous sulphate01 g
Distilled water	1000 ml.

Each medium will be prepared by mixing the requisite amounts of the ingredients. It will be sterilized in autoclave at 15 lbs. pressure for 20 -35 mts. The sterilized medium will be poured in sterilized petri dishes and will be inoculated with equal quantity of inoculum.

The growth of the pathogen on the semi solid medium will be measured in centimeters. While, the growth of the pathogen in liquid medium will be measured on the basis of dry weight of mycelium after 21 days of growth. The contents will be filtered through Whatman filter paper No.1 previously dried and weighed. Filter papers containing mycelium will be dried at 60°C for 24 hours in crucibles, after they cool the weight will be taken. All the experiments will be repeated thrice and 5 replications will be used in each case.

Effect of different carbon and nitrogen sources-

Carbon and Nitrogen requirements will be studied

by replacing glucose and asparagine of the basal medium with different quantities of C & N compounds. The quantity of carbon sources added will be determined on the basis of their molecular formulae, so as to contain an equivalent amount of carbon as present in 10 gm. of the glucose in basal solution. Similarly quantity of nitrogen compounds will be added to the basal solution on the basis of nitrogen present in 1 gm. of asparagine.

Carbon sources to be used:- Xylose, mannose, galactose, Arabinose, Fructose, glucose, sucrose, maltose, starch, mannitol, sorbitol.

Nitrogen sources to be used:- Ammonium sulphate, Ammonium nitrate, ammonium chloride, Potassium nitrate, sodium nitrate, urea, asparagine, alanine.

Effect of mineral elements:-

Effect of mineral elements on the growth of mycelium will be studied by following the method of Steinberg (1946) with some modifications. $MgSO_4$ will be substituted by $(NH_4)_2SO_4$, $CaCl_2$ by NH_4Cl and $NaNO_3$ by NH_4NO_3 respectively.

Effect of temperature:-

150 ml. Erlenmeyer flasks, filled with 50 ml. of sterilized liquid basal medium will be inoculated with an equal amount of fresh inoculum. The flasks will be incubated at 9 different temperatures maintained at intervals of $5^{\circ}C$ i.e., 0° , 5° , 10° , 15° , 20° , 25° , 30° , 35° & $40^{\circ}C$.

Effect of H-ion concentrations on the growth of the Pathogen:-

Erlenmeyer flasks of 150 ml. capacity containing sterilized liquid medium having different P^H will be inoculated with equal amount of mycelium taken out from the fungal colonies. These flasks will be incubated for 15 days at room temperature. The P^H value of the medium will be adjusted by Beckman's P^H meter using N-NaOH and HCl before autocleaving. 12 levels of P^H i.e. 3.5, 4.0, 4.5, 5.0, 5.5, 6.0, 6.5, 7.0, 7.5, 8.0, 8.5 & 9.0 will be tried.

Biochemical Studies:-

Under this head the post-infection changes caused by the pathogen will be studied. Healthy fruits and vegetables inoculated and together with uninoculated fruits and vegetables shall be analysed for sugars, aminoacids, ascorbic acid, phenolics, organic acids and enzymes.

For these studies 5g pulp in the case of AA's 50g pulp from the healthy as well as inoculated fruits and vegetables will be macerated in ground - glass homogenisers containing acid-washed sand and 20 ml. of 80% ethanol. The macerated pulp will be boiled in 20 ml. of 80% ethanol on a water bath. Care will be taken to obtain even the last traces of organic compounds. The solution thus obtained will be kept at a lower temperature, the supernatant thus obtained will be centrifuged at 2,000 rpm for 30 mts. These studies will be initiated after 3 days of inoculation and will be continued upto 12 day of inoculation.

Sugars:-

One directional chromatography technique (Block et al 1958) will be followed for the estimation of sugars. The solvent system used will be n-butanol:pyridine: water in the ratio of 6:4:3. A straight line curve with a pencil will be drawn on the Whatman filter paper No.1, on this line a diluted extract will be spotted in triplicate by means of a self filling Lane de pipette. In order to keep the spots to a minimum size, they will be dried with the help of an air blower. The chromatograms shall be run for 18 hrs. in the saturated chromatographic chamber with the solvent. The chromatogram will be taken out of the chamber and dried at room temperature till the solvent evaporates. The developing reagent (5 vol. of 4% aniline + 5 vol. of diphenylamine + 1 vol. of orthophosphoric acid) will then be sprayed after this the chromatograms shall be dried at room temperature and then heated at 110°C for 15 mts.

Quantitative estimation:-

Spots from unsprayed chromatograms will be elused with hot distilled water. By the help of phenol - sulphuric acid method the sugars shall be estimated by using Bausch & Lomb Spectronic - 20 spectrophotometer at wave length of 440 m.μ. The quantities shall be determined by referring to a standard curve prepared with authentic samples of sugars.

Free amino acids & organic acids:-

For the estimation of amino acids 2 dimensional ascending technique shall be used. Samples each measuring 30 ml. will be spotted on Whatman paper No.1. The solvents used in this case will be n butanol: acetone: H_2O in the ratio of 120: 30 : 5 followed by liquid phenol: water: ammonia in the ratio of 180:20:1. In each direction the chromatograms shall be run for 18 hrs. and then dried at room temperature after each run. Ninhydrin solution (5%) shall be used as a developer. The sprayed chromatograms shall be left at room temperature for an hour and later heated at $45^{\circ}C$ for 10 mts. The coloured spots will be removed and will be read in Bausch & Lomb Spectronic - 20 spectrophotometer at 540 n.m. While proline and asparagine will be read at 440 n.m. and 530 n.m. respectively. The quantities shall be determined by reference to a standard curve prepared with authentic samples of amino acids.

For the estimation of organic acid Uni-directional ascending paper chromatography shall be used. The solvents used in this case shall be n-butanol: formic acid: H_2O in the ratio of 4:1:5. The chromatograms shall be run for 12 hrs. and sprayed with aniline - xylose reagent then heated at $130^{\circ}C$ for 20 mts. The spots will be eluted with 70% ethanol. Optical density will be measured at 365 n.m. Quantitative analysis shall be carried out with the help of authentic samples of organic acids.

Enzymes:-

Pulp (10g) from healthy as well as diseased specimens will be macerated in ground glass homogenizer with 20 ml. of 0.2N NaCl solution in distilled water. The homogenate will be filtered through several layers of muslin cloth and squeezed. The extract will then be centrifuged at 400 r.p.m. for 20 mts. and the final volume will be made 25 ml. with 0.2 N NaCl solution. This test solution will be used for all enzymatic assay.

1. Pectolytic enzymes:-

(A) Pectin - methyl esterases:-

This was carried out by continuous titration method as described by Kertesz (1951), followed by Goodenough & Maw (1974).

For this a solution of 1.2% citrus pectin with P^H adjusted at 5.5 with the help of 1N NaOH will be used as a substrate. 30 ml. of this substrate will be taken in a beaker and to this 4.5 ml. of enzyme preparations will be added. P^H of the reaction mixture will be noted at experimental time zero. After 120 mins. P^H will again be recorded. The original P^H will be recovered by titrating the mixture with .025 N NaOH. The amount of NaOH consumed to get original P^H will be recorded.

The results obtained will be expressed in terms of micro equivalents of methoxy groups released/ml./120 min by the help of the formula suggested by Smith (1958).

Microequivalents of methoxy groups released/ml. =
 .025N NaOH consumed to get original P^H X .775
 4.5

(B) Polygalacturonase (PG):-

With the help of Viscometric method followed by Hall and Wood (1974) & Bisen & Agarwal (1980) PG will be assayed. Reaction mixture consisting of 1.2% solution of sodium polypectate, 2 ml. of .2 M. will be prepared McIlvain buffer (P^H 4.6), 1 ml. of distilled H₂O and 1 ml. of enzyme preparations. Efflux time will be noted after a definite time interval. The enzyme activity will be expressed in terms of % loss in viscosity and relative enzyme activity, the formula of Cappeline (1966) will be applied for the same.

$$1. \% \text{ loss in viscosity} = \frac{E_{T_0} - E_{T_t}}{E_{T_0} - E_{T_v}} \times 100$$

Where E_{T₀} = Efflux time in seconds for control

E_{T_t} = " " " " " reaction mixture at time t

E_{T_v} = " " " seconds for water.

2. Relative enzy activity (REA)

$$= \frac{100}{t \text{ at } V_{50}}$$

Where t at V₅₀ = Time in mins. required to reduce the viscosity by 50%.

Reaction mixture containing distilled H₂O or boiled enzyme preparations in-stead of active enzy preparations will serve as control.

(C) Polyacetyl - galacturonase (PMG)₁-

PMG will also be assayed by the same Viscometric method which was used by Bell et al (1955), Keen & Horton (1966). The reaction mixture will consist of the following composition.

Citrus pectin (1.2%)	2 ml.
.2 M Mollvain buffer (5 p^H)	2 ml.
Distilled H ₂ O	1 ml.
Enzyme preparation	1 ml.

2. Cellulolytic enzymes₁-

Cellulase (CX)₁- CX enzyme will also be assayed by viscometrical method (Hancock et al 1964 a, Hall & Wood 1974). The composition of the reaction mixture will be as follows:

.5% Carboxymethyl cellulose	2 ml.
.2 M Mollvain buffer (p^H 5.5)	2 ml.
Distilled H ₂ O	1 ml.
Enzy preparation	1 ml.

The results will be expressed in % loss in viscosity and relative enzyme activity.

Effect of fungicides, phenols & antibiotics on the enzyme activity₁-

Effect of various chemicals on the activity of PG, PMG & CX will be investigated by using the same viscometric method. The reaction mixtures will be the same as used previously only in place of distilled water 1 ml. of effective substance of desired concentration will be added. Different concentrations of the following substances will be added:-

(a) Antibiotics:-

	Concentration (ppm)		
(a) Tetracycline	100	500	1000
(b) Nystatin	"	"	"
(c) Chloramphenicol	"	"	"

(b) Fungicides:-

	Concentration (ppm)		
Copper oxychloride	100	500	1000
Brassicol	"	"	"
Morestan	"	"	"

(c) Phenols:-

B-naphthol	"	"	"
Phloroglucinol	"	"	"
Dinitrophenol	"	"	"

Effect of different packing systems:-

To determine the effect of different packing systems on the development of post harvest diseases, the freshly obtained fruits and vegetables will be packed in wooden crates with different kinds of packing material such as wood-straw, rice straw, hay, plastic cuttings, paper cuttings. The fruits before packing will be rinsed with different detergents and sterilising agents. Different combinations of packing material and sterilizing/detergent agents will be tried. After every 3 days the packing will be opened and the intensity of various diseases development will be determined.

REFERENCES

REFERENCES

- Abdel - Sattar, M.A.; M.M. Satour and A.A. El-Shohedi
(1979): Ceratocystis fruit rot disease of banana in
Egypt. Agricultural Research Review, (1977), 55 (2):
79-85 Pl. Path Res. Inst. Agri. Res. Cent. Egypt.
- Agarwal, R.K. and V.C. Sharma (1968): Storage of disease of
apple fruit. Indian Phytopath. 21: 294-298.
- Akamine, E.K. (1977): Tropical and Subtropical crops in
Hawaii. CSIRO PD D Res. Q. 37 (1): 13-20.
- Akamine, E.K. and T. Soo (1969): Effects of controlled at-
mosphere storage of fresh papayas (Carica papaya L.
var Solo) with special reference to shelf-life ex-
tension of fumigated fruits. Hawaii Agric. Exp. Sta.
Res. Bull 144, 27 pp.
- Ali, S.S. (1976): Changes in sugar content of mosambi fruits
infected with Botryodiplodia theobromae Pat. Acta
Bot. Indica, 4:63-65.
- Ali, S.S. and A. Shukla (1981): Pathogenesis of Rhizopus
oryzae Went & Geerlings in fruit rot disease of
brinjal (Solanum melongena L.) Current Science (1981)
30 (5):232-233. Ravishankar University, Raipur, India.
- Alvarez- Garcia, Luis (1972): Thielaviopsis rot of bananas
J. Agric Univ. PR, 55 (2): 261-262.

Aragaki, M. and J.Y. Uchida (1979): A new papaya fruit rot in Hawaii caused by Phytophthora capsici. Plant Disease Reporter. (1978) 62 (9): 765-768. Univ. Hawaii Honolulu USA.

Artschwager, E. (1927): Wound periderm formation in potato as affected by temperature and humidity. J. Agric. Res. 17: 137-52.

Ashby, B.H. (1970): Protecting perishable foods during transport by motor truck. US Dep. Agric. Handb. 105, 141 pp

ASHRAE (1974): ASHRAE Guide and Data Book- Applications for 1974 Am. Soc. Heating, Refrig. Air-Cond. Eng. New York 1974 ed.

Aulakh, K.S.; R.K. Grover and S. Malhotra (1970a): Utilization of free aminoacids in vivo and in vitro by isolates of Phoma destructiva Plowr. Phytopath. Medit. 9: 8-12.

Aulakh, K.S.; R.K. Grover and S. Malhotra (1970b): Changes in vitamin C & Carbohydrate content of tomato fruits inoculations with isolates of Phoma destructiva Phytopath Medit. 9: 91-94.

Avarez- Garcia, Luisa & Justo Lopez-Garcia (1972): Gummosis, die-back and fruit rot disease of mango (Mangifera indica L.) caused by Physalospora rhodina (B & C) Cke in Puerto Rico. J. Agric. Univ. PR. 55 (4): 435-450.

- Bompeix, G.; M. Coeffic and P. Greffier (1980): Control of peach rots due to Monilia spp Batrachia sp & Rhizopus sp. (1979) 34 (6): 423-430. Dep Pl. Path Univ. P & M Curie, Paris 6, France.
- Borecka, H. (1979): Fungi of the genus Penicillium on apples & pears during the storage period. Acta Agrobotanica. (1977) 30 (2): 213-227. Res. Inst. Pomology, Skierniewice, Poland.
- Bourne, M.C. (1976): Proposed definition of post harvest feed loss. Proc. Natl. Feed loss conf. Boise, Idaho, pp 129-30.
- Braslauska, O. (1978): The pathogen of dry rot of potato - Fusarium solani (Mart) Sacc in Central Slovakia (1977) 13 (2): 97-100. Cent. Agric. Testing Inst., Zilina, Czechoslovakia.
- Brodrick, H.T. and G.C.A. Vander Westhuisen (1977): Hendersonia groberrima, the cause of soft brown rot of mango in South Africa. Phytophylactica 8 (1): 13-16.
- Brook, P.J. (1978): Glomerella singulata & bitter rot of apple. New Zealand Journal of Agricultural Research (1977) 20 (4): 547-55. Plant. Dis. Div. DSIR AUCKLAND, New Zealand.
- Byrde, R.J.W. and H.J. Willette (1977): The brown rot fungi of fruit. Their biology & control. Oxford U.K. Pergamon Press (1977) 171 pp.

- Bhargava, S.N.; P.N. Shukla and Narendra Singh (1981): Some studies on Botryodiplodia rot of apple proceeding of the National Academy of Sciences, India B. (1979) 49 (1): 43-45. Allahabad Univ., Allahabad, India.
- Bhatia, I.S.; D.S. Uppal & K.L. Bajaj (1972): Studies on phenolic contents of resistant and susceptible varieties of tomato, (L. esculentum) in relation to early blight disease Indian Phytopath. 25 : 231-235.
- Bhutani, B.C. (1946): Vitamin contents of the mango. Punjab Fruit J. 10: 112-115.
- Bisen, P.S. and G.P. Agarwal (1980): In vitro production of pectolytic enzymes by Aspergillus niger van Tiegh causing soft rot in apples. Phytopathologische Zeitschrift, (1980), 97 (4): 317-326 Univ. Jabalpur India.
- Block, R.J., E.L. Durrum and G. Zweig (1958): A manual of paper chromatography & paper electrophoresis Academic Press, Inc., Publishers New York.
- Bolay, A. (1978): Phytophthora syringae, causal agent of a serious rot of stored apples. Revue Suisse de Viticulture d' Arboriculture et d' Horticulture, (1977), 9 (4): 161-169.
- Bolkan, H.A., W.R.D. Ribeiro and O.C.D.E. Almeida (1981): Pathogenicity of Cylindrocarpum clavatum causing potato tuber rot. Plant Disease, (1981) 65 (1): 47-49 Univ. Brasilia, Brazil.
- Babovic, M.; M. Perisic.; S. Markovic; S. Stojanovic, Z. Pente-

- lic. (1980): Investigation on rot of apple fruits in cold store house Zastita Biliana (1979) 30 (147): 83-87. Vas. Agric. Belgrade, Zemun, Yugoslavia.
- Bai, D.H. (1979): Survey on the fruit rot occurrence and damages of shipping mandarin. Korean Journal of Plant Protection. (1977) 16 (4): 245-247. Daihan Agric. Science Inst. Korea.
- Barts, J.A. (1981): Causes of post harvest losses in a Florida tomato shipment Plant Disease. (1980) 64 (10): 934-937 Univ. Florida, Gainesville USA.
- Bell, T.A., J.L. Etchells & T.D. Jones (1955): A Method for testing cucumber salt-stock brine for softening activity. U.S. Dept. Agr. ARS 72: 5-15 p.
- Bhargava S.N. and A.P. Singh (1980): Aspergillus rot of mango its control. Indian Journal of Horticulture. (1975) 32 (3/4): 190. Univ. Allahabad India.
- Bhargava, S.N.; D.N. Shukla and Narendra Singh (1981): Post harvest diseases of cherry. Proceedings of the National Academy of Sciences. India B. (1978) 48 (3): 168. Allahabad Univ., Allahabad, India.
- Capellini, R.A. (1966): Growth and polygalacturonase production by Rhizopus stolonifer. Phytopath 56: 734-737.
- Carroll, R.B. and M. Sasser (1974): A outbreak of potato pink rot in Delaware. Plant Dis. Rep. 58 (5): 443-445.

- Chahal, A.S. & R.K. Grover (1972): Changes in ascorbic acid, aminoacids and carbohydrates content of chilli fruits with Cheeseophora cucurbitarum. Indian Phytopath. 25: 257-260.
- Chandra, S. & R.N. Tandon (1963): Studies on rotting of apple, banana, lemon and tomato fruits by fungi. Phyton, 26: 41-47.
- Chary, S.J. & B. Pradeep Kumar & S.M. Reddy (1980): Hitherto unrecorded post harvest disease of tomato. Indian Phytopathological society, (1980) 33 (4): 624-625.
- Chaurasia S.C. (1980): A new fruit rot disease of India (Citrullus vulgaris Schard var-fistulosus Stock) Indian Phytopath. 33 (2): 310.
- Chemulu, V.V. & D.P. Thakur (1968): Survey of storage & market disease of certain fruits and vegetables in Delhi and control of soft rot due to Rhizopus Sp. Indian Phytopath. 21: 133-143.
- Chohan, J.S. & T. Chend (1981): Phoma glomerata a new pathogen on pears (Pyrus communis). Transactions of the British Mycological Society, (1980) 75 (3): 501-511. Punjab Agric. Univ. Ludhiana, India.
- Chowdhury, S.R. and S.K. Hasija (1981): Pathological studies on Phomopsis vexans causing soft rot of brinjal fruits Indian Phytopathology, (1979. publ. 1980) 32 (3) 493-496. Univ. Jabalpur M.P. India.

- Datar, V.V. (1981): Brinjal fruit rot caused by Fusarium moniliforme Sheld- a new record from India (Gibberella fujikuroi on egg plant) Current Science, (1980) 49 (14): 555. Fruit. Res. Store, Himayat Bagh, Aurangabad, India.
- Dennis, C. (1981): Post harvest spoilage of strawberries ARC Research Review, (1978) 4 (2): 38-42. Food Res. Inst. Colney Lane, Norwich, U.K.
- D'ercolo, N. (1981): Phytophthora nicotianae var nicotianae casual agent of tomato rot under cover. Informatore Fitopatologico, (1980) 30 (11/12): 21-24. Institute di Patologia Vegetale Univ. Bologna Italy.
- D'ercolo, N. and P. Flori (1978): A case of peach rot due to Phytophthora syringae (Kleb) Kleb. Informatore Fitopatologico, (1978) 28 (5): 19-21. Institute di Patologia Vegetale, Univ. Bologna Italy.
- Devey, D.H. (1977): Controlled atmospheres for the storage & transport of perishable agricultural commodities Mich. State Univ. Hortic. Rept. 28 301 pp.
- Devey, D.H. R.C. Herner, D.C. Dilley (1969): Controlled atmospheres for the storage and transport of horticultural crops. Mich. State Univ. Hortic. Rept. 9, 153 pp.
- Dhingra, R. and R.S. Mehrotra (1980): A few unrecorded post-harvest diseases of fruits and vegetables. Indian Phytopath (1980) 33 (3): 475-476.
- Dhingra, R., R.S. Mehrotra and K.R. Anuja (1981a): An unrecorded fruit rot of orange (Citrus reticulata Blanco) (Acremonium

- synonym) Current Science, (1980) 49 (8): 329. Kurukshetra Univ. India.
- Bhingra R., K.S. Mehrotra and K.R. Anje (1981b): A new post harvest disease of Annona squamosa L. (Glomerella cinerata). Current Science (1980) 49 (12): 477-478. Kurukshetra Univ. Harayana India.
- Eckert, J.W. and N.F. Sommer (1967): Control of diseases of fruits and vegetables by post harvest treatment. Ann. Rev. Phytopathol. 5: 391-432.
- Eiche M.M. (1981): (Pathogens of storage rots of apple in Latvia) Inst. Pl. Prot. Baltic Branch, USSR from Referativnyi Zhurnal, Fitopatologiya (1980) 11-79. (193).
- El-Khadem, M. and M.K. El-Kassas (1978): Acremonia atra (Corda) Sacc isolated from dry rotted potato tubers in Egypt. Egyptian Journal of Phytopathology, (1975) Tanta Univ. Kafr El-Sheikh Egypt.
- Feld, S.J., J.A. Manges, J.E. Pehrson (1979): Brown rot of Citrus a review of the disease. Citrographa (1979) 64 (5): 101-106 Univ. California, Riverside, Calif, U.S.A.
- Ferrin, D.M. and D.C. Ramadell (1978): Ascospore dispersal & infection of grapes by Guignardia bidwellii, the causal agent of grape black rot disease. Phytopathology (1977) 67 (12): 1501-1505 Michigan State Univ. East Lansing USA.
- Garcha, Harnek, Singh and Vapinder Singh (1980): Post harvest disease of fruits in Punjab Indian Phytopathology, 1980 22 (1): 42-47.

- Garg, P.K. and M.N. Gupta (1981): A Fusarium rot of tomato fruits
Indian Phytopathology (1979, publ. 1980) 32 (2): 332-333.
Agra College, Agra.
- Genry, J.P. and R. Guillou (1966): Fogespray humidification in cold
storage Am. Soc. Heat Refrig. Air-Cond. Eng. J. 8 (10):
80-83.
- Gentry, J.P.; F.G. Mitchell and N.F. Sommer (1965): Engineering
and quality aspects of deciduous fruits packed by volume-
filling and hand-placing methods. Trans. Am. Soc. Agric.
Eng. 8: 584-585, 89.
- Georgieva, M. and S. Kotev (1979a): Rot of cucumber fruits during
transportation and its control. From abstracts of Bulgarian
Scientific Literature A 22 (4): 821.
- Georgieva, M. and S. Kotev (1979b): Penicillium rot of garlic *Sine*
Plesenyasvane (*Penicillium guienie*) *Pe Ches'na Gradinarska*
i Lovarska Nauka (1977) 14 (8): 44-51, Kolarov Higher Inst.
Agric. Plovdiv, Bulgaria.
- Georgieva, M. and S. Kotev (1979c): The Pathogens of garlic rot
during storage from abstracts of Bulgarian Scientific
Literature A 22 (4): 822.
- Gerhardt, F. (1955): Use of film box liners to extend storage life
of pears and apples US Dep. Agric Circ 965 28pp.
- Ghosh, A.K. and R.W. Tandon (1965a): Formation of a new oligosac-
charides in mango fruits under pathogenesis. Curr. Sci.
34: 465.

- Ghosh, A.K.; R.N. Tandon; S.N. Bhargava and M.P. Srivastava (1964):
Vitamin C content of guava fruits after fungal infection.
Naturwissenschaften 52: 478.
- Glaser, T. (1979): Black rot of carrot. 98 71-85. Inst. Pl. Prot.
Poznan. Poland.
- Columbie, C. (1964): Maintaining quality of farm crops. In year
book of Agriculture, ed. A. Stefferud, pp 291-302
Washington DC, USDA. 592 pp.
- Goodenough, P.W. and G.A. Maw (1974): Studies on the root rotting
fungus Pyrenochaeta lycopersiae characterization of extra-
cellular enzymes with degrade pectic compound and factors
affecting the synthesis of these enzymes. Physiol. Pl.
Pathol. 4: 57-62.
- Granger, K. and A.S. Horne (1924): A method of inoculating the
apples. Ann Bot (N.S.) 38: 212.
- Grierson, W. (1969): Some random thought on CA research Hicha
State Univ. Hortia. Rept. 9, pp 77-79.
- Guillou, R. (1960): Coolers for fruits and vegetables Calif
Agric. Exp. Stn. Bull 773. 66pp.
- Gupta, M.L.; L.N. Dubey and V. Nath (1978): Storage rot of
potato caused by Aspergillus flavus Link (1935) Geobion
(1977); 4 (2) 80. Univ. Gorakhpur India.
- Gupta, P.C. and R.L. Madan (1979): Fruit rot disease of ber
(Zizyphus mauritiana L.) from Haryana. Indian Phytopa-
thology (1977, publ. 1978); 20 (4): 554-555. Haryana
Agric. Univ. Hissar India

- Gupta V.K. and K.D. Verma (1980): Dry rot of cherry fruit caused by Alternaria dendritica Indian Phytopathology (1978, publ. 1979); 31 (4): 512-513. Himachal Pradesh Univ., Coll. Agric., Solan, India.
- Gusev S.A. (1981): The storage conditions and diseases of potato Zashchita Rastenii (1980); 10: 54-56. NIIKKh, USSR.
- Haegermark, U. (1980): White rot (Sclerotium cepivorum Berk) - a parasitic fungus on onion in the island of Oland. 43 (5/6): 102-108.
- Hall, J.A. and R.K.S. Wood (1974): Permeability change in tissues and other effects of cell separating solutions from soft rot caused by Corticium pratense & Erwinia Atroseptica Ann Bot 38: 129-140.
- Hancock, J.G.; R.L. Millar and J.W. Lorbeer (1964a): Pectolytic and cellulolytic enzymes produced by Botrytis allii, B. cinerea and B. squamosa 'in vitro' and 'in vivo' Phytopath 54: 928-931.
- Hardenburg, R.E. (1971): Effects of in-package and environment on keeping quality of fruits and vegetables. Hort-Science 6: 198-201.
- Hardenburg, R.E. (1973): Use of plastic films in maintaining quality of fresh fruits and vegetables during storage and marketing. Am. Soc. Heat Reftig. Air-Cond. Eng. Symp 1973, pp. 19-29.

- Hardenburg, R.E. (1974): Benomyl and thiebendazole treatments reduce decay of apples in polyethylene consumer bags. J. Am. Soc. Hort. Sci. 99: 236-38.
- Harvey, J.M. (1978): Reduction of losses in fresh market fruits and vegetables. Ann Rev Phytopathol. (1978); 16: 321-41.
- Harvey, J.M.; H.M. Cousey, H.C.M. Porter and F.M. Porter (1966): Air transport of California strawberries US Dep Agric Mark Res. Rept. 751. 12 pp.
- Harvey, J.M. and M. Vota (1977): Table grapes and refrigeration: Modified atmospheres, in particular the influence of SO₂ Bull. Inst. Intern. Frigid (Paris) in press.
- Hasijs, S.K. and H.C. Agarwal (1977): Pathological studies on Trichothecium roseum causing rot of apple and plum fruits. Indian Phytopath. 30 (2): 254-255.
- Hasijs, S.K. and S. Datta (1981): Pathological studies on Phoma destructiva causing rot of tomato fruits. Indian Phytopathology (1979, publ. 1980); 32 (2): 327-328 Univ. Jabalpur, India.
- Haware, M.P. and L.K. Joshi (1976): Studies on soft rot of ginger from M.P. Indian Phytopathol. 27 (2): 158-161.
- Howard, Charles M. (1972): A strawberry fruit rot caused by Colletotrichum fragariae. Phytopathology 62 (6): 600-602.
- Howard, C.M. & L.E. Albregts (1973): A strawberry fruit rot caused by Hendrophoma obscurans. Phytopathology 63 (3): 419-421.

- Howard C.M. & E.E. Albrechts (1974a): A strawberry fruit rot caused by Pestalotia longisetula. Phytopathology 63 (7): 862-863.
- Howard C.M. & E.E. Albrechts (1974b): A strawberry fruit rot caused by Alternaria tenuissima. Phytopathology 63 (7): 938-939.
- Ikediugwu, P.M.O. (1981): Corticium rolfeii a fruit rot of Citrus limon in the field in Nigeria. Transactions of the British Mycological Society. (1980) 75 (2): 316-319. Univ. Benin, Benin City, Nigeria.
- Ikediugwu, P.M.O. & W.O. Ogilewa (1979): Fruit rot of Citrus limon in Nigeria caused by Fusarium solani. Transactions of the British Mycological Society (1978): 71 (2): 209-213 Univ. Benin, Nigeria.
- Jahn, N. and U. Burth (1981): On the occurrence of parasitic storage rots of apple. Nachrichtenblatt für den Pflanzenschutz in der DDR. (1981) 35 (4): 76-79.
- Jamaluddin (1979): Organic acid composition of peach fruits during infection by Aspergillus flavus. Indian Phytopathology 32 (3): 490.
- Jamaluddin (1981): Gliocladium rot of fruits of Phyllanthus emblica. Proceedings of the National Academy of Sciences B. (1978) 48 (1): 62. Allahabad Univ. Allahabad India.
- Jamaluddin, M.P. Tandon and R.N. Tandon (1978): Rot of fruits litchi (Litchi chinensis) in marketing processes (Geotrichum candidum) Indian Phytopathology (1975, publ. 1976)

28 (4): 530-531. Allahabad Univ., India.

Jamaluddin, M.P. Tandon and R.M. Tandon (1979): A fruit rot of 'Aonla' (Phyllanthus emblica L.) caused by Phoma. Proceedings of the National Academy of Sciences, India B. (1975) 45 (1): 75-77 Univ. Allahabad, India.

Janke, C. and A. Zett (1981): Occurrence of Phoma dry rot on stored potatoes in the GDR. Archiv für Phytopathologie und Pflanzenschutz (1980); 16 (5): 325-332 Humboldt Univ., Berlin, GDR.

Jones, A.L. and W.G. Fields (1972): An apple fruit rot caused by a Hormiscium sp. Plant Dis Rep. 56 (4): 275.

Kable, P.F. (1973): Brown rot of stone fruits on Murrumbidgee Irrigation Areas: III. Aust J Agric Res. 23 (6): 1035-1044.

Kamal: R.P. Singh and K.S. Bhargava (1978): Sour rot of lemon new to India. Indian Phytopathological Society; 31 (1): 76-77.

Kanaujia R.S. (1979): Fruit rot of mandarin orange Citrus reticulata. Indian Phytopathology 32 (3): 450.

Kanaujia R.S. (1981a): A new Aspergillus storage rot of mandarin orange in India (Aspergillus aculeatus) Indian Phytopathology (1979 publ 1980); 32 (4) 620-621.

Kanaujia R.S. (1981b): Fruit rot of mandarin orange (Trichoderma viride). Indian Phytopathology (1979, publ. 1980); 32 (3) 450

- Kao, C.W. and L.S. Lou (1981): Strawberry fruit rot caused by Phytophthora cactorum and P. citrochthona. Plant Protection Bulletin Taiwan (1979); 21 (2): 239-243. Pl. Prot. Cent. Wufeng, Taiwan.
- Kapeor, I.J. (1968): Pathological studies of the fruits rot diseases. D. Phil. Thesis, Allahabad Univ., Allahabad, India.
- Kapeor, I.J. and R.N. Tandon (1967): Formation of a new oligosaccharide in guava fruits under pathogenesis Curr. Sci. 38: 444.
- Kapeor, I.J. and R.N. Tandon (1969): Post infection changes in ascorbic acid content of tomato fruit caused by Drechslera australiensis. Curr. Sci. 38: 397-398.
- Kaul, J.L. and R.L. Munjal (1981): Post infection biochemical changes in apple fruits due to rot causing fungal pathogens. Gartenbauwissenschaften (1980); 45 (4): 185-187. Himachal Pradesh Agric Univ. Solan, India.
- Keen, N.T. and J.C. Horton (1966): The Polygalacturonase of Pyrenopeziza terrestris Phytopath. 56: 603-609.
- Kertesz, Z.I. (1951): The pectic substances Interscience publishers New York 628 p.
- Khanna, K.K. and S. Chandra (1979a): Studies on storage disease of fruits and vegetables III Factors affecting storage rot of lime caused by Aspergillus niger. Proceedings of the National Academy of Sciences India B. (1975) 45 (2): 109-110, Univ. Allahabad U.P. India.

- Khanna, K.K. and S. Chandra (1979 b) Studies on storage diseases of fruits and vegetables II, Factors affecting banana fruit rot caused by Fusarium moniliforme and Fusarium roseum. Proceedings of the National Academy of Sciences India B. (1975); 45 (2): 105-108 Univ. Allahabad, India.
- Kogan, E.D. (1979): The pathogen of black rot of tomato fruits in Moldavia. From Referativnyi Zhurnal (1977); 2 (55):1248.
- Kritagyan S.P. Sahu and S.P. Singh (1980): Fruit rot of pointed gourd (Trichosanthes dioica) in Bihar, Indian Phytopath. 33 (2): 308-309.
- Kumar, S.; and M.P. Tandon (1979): A new fruit rot of carissa Myrothecium poridum on C. carandas Indian Phytopathology (1978); 31 (1): 105 University Allahabad India.
- Kumbhare, G.B. and K.G. Choudhari (1980): Storage decay of sweet orange and control measures. Journal of Maharashtra Agricultural Universities (1979); 4 (2): 230-231.
- Kunimoto, R.K.; P.J. Ito and W.H. Ko (1977): Mucor rot of guava fruits caused by Mucor hiemalis. Tropical Agriculture (1977); 54 (2): 185-187. Coll. Trop. Agric. Univ. Hawaii Hilo Hawaii USA.
- Lal, B.; R.N. Rai and A. Arya (1980): Colletotrichum rot of papaya National Academy Science Letters (1979); 3 (5): 174. Allahabad Univ. India.

- Langerfield E. (1979): Fusarium coeruleum (Lib.) Sacc as the casual agent of storage rots on potato tubers. Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft Berlin Dahlem (1978) 184, 81 pp.
- Langerfield (1980): Phoma eupyrena Sacc causing rot on Potato tubers. Nachrichtenblatt des Deutschen Pflanzenschutzdienstes (1979); 31 (7): 11.
- Lentican, M.T. and T.H. Quisio (1978): Pathogenicity and cultural characteristics of Botryodiplodia spp causing fruit rots. Philippine Phytopathology (1976); 12 (1/2): 66-74. Univ. Philippines Los Banos College Laguna, Philippines.
- Laxminarayan, P. and S.M. Reddy (1978): A new post harvest disease of mango. Indian Phytopathology (1975, publ 1976); 28 (4): 529-530. Post graduate cent, Walangal A.P. India.
- Laxminarayan, P. and S.M. Reddy (1980): Some new post harvest diseases of chillies. Indian Journal of Mycology and Plant Pathology, (1978), publ. 1979); 8 (2): 202-203. Kakatiya Univ., Vidyaranyaपुरi India.
- Lee, C.U. (1978): Effect of post harvest temperature on potato piece rot in relation to suberin and periderm development. Korean Journal of Plant Protection (1977) 16 (1) 55-63. Seoul Natn Unit. Suwon, S. Korea.
- Lents, C.P. and L. van den Berg (1971): Study of factors affecting temperature, relative humidity, and moisture loss in fresh fruit and vegetable storages. J. Inst. Can. Technol. Aliment; 4, 46-53.

- Lipton, W.J. (1967): Market quality and rate of respiration of head lettuce held in low-oxygen atmospheres. US. Dep. Agric. Mark. Res. Rept. 777, 9pp.
- Lockhart, G.L. (1969): Effect of CA storage on storage rot pathogens. Mich. State Univ. Hortis. Rept. 9, pp 113-21.
- Lopkova, K.V. (1976): Phytophthora erythroseptica Pethybr-pathogen of pink watery rot of potatoes. Mikol Fitopatol; 8 (6): 524-525.
- Lusas, H.T.; Antonia, B. Carvalho and Maria-C. Barrioli (1975): Pear fruit rot caused by Coremiella subispora. Trans Br Mycol Soc; 62 (1): 206-208.
- Lutz, J.M. and R.E. Heidenburg (1968): The commercial storage of fruits, vegetables and florist and nursery stocks. US Dep Agric Handb. 66, 94 pp.
- MacLeod, R.F., A.A. Kader and L.L. Morris (1976): Damage to fresh market tomatoes can be reduced Calif. Agris. 20 (12): 10-12.
- Maiti, S; H.S. Shanbi and S.P. Ghosh (1981): A new fruit rot on plum (Prunus domestica) Glomerella singulata Indian Journal of Mycology and Plant Pathology, (1979, publ. 1980); 2 (2): 249. Indian Council Agric. Res. North Eastern Hill Reg. Shillong India.
- Majumdar, G. and V.V. Modi (1981): Spoilage of mango by Aspergillus flavus Current Science, (1980) 49 (21): 821-822. Ms University Baroda India.

- Nathur, S.B. and B.L. Nathur (1958): Indian Phytopath, 12: 66-69.
- Nathur, S.B. and V.K. Agarwal (1974): A Pestalotia sp causing fruit rot in apples. Prissia, 10 (1/2): 57-60.
- Narte, M. and F. Tazburi (1980): A case of dry rot of potato due to Fusarium trichotecoides Wall. Fitopatologia, (1979) 22 (11/12): 11-14.
- Matsuzaki, M., H. Kan and A. Kiso (1981): Phytophthora rot of strawberry in Kyushu Japan, Annals of the Phytopathological Society of Japan (1980) 46 (2): 179-184.
- Mac cullen W.D. and William B. Howitt (1973): Early Botrytis rot of grapes: Time of infection of latency of Botrytis cinerea Pers in Vitis vinifera L. Phytopathology, 63 (9): 1151-1157.
- Mc Combs C.L. and N.N. Winstead (1964): Changes in sugars and AAs of cucumber fruits infected with Pythium aphanidermum. Phytopath, 54 (1): 233-234.
- Mc Intosh, D.L. and S.W. Porritt (1958): The effect of storing apples in polythene box liners on the development of Bull's Eye Rot, caused by Nectria perennans. Plant Dis Repr., 42 97-99.
- Mehta, P., K.M. Vyas and S.B. Saxena (1977): Pathological studies on fruit rot of tomato caused by Alternaria solani and A. tenuis. Indian Phytopath, 28: 247-252.
- Mehta, P., K.M. Vyas and S.B. Saxena (1977): Metabolic changes

during pathogenesis of fruit rot diseases of tomato.

Indian Phytopath. 28 253-255, 1975(1976).

Michail, Samir, H., A. Mohamed, Abd- El-Rehim, Husain Elarosi
and Enayat A. Khairy (1972): Dry rot of garlic cloves
in U.A.R. (Egypt), Phytopathol. Mediter 10 (2): 202-205.

Nicosia, R.S. and L.L. Ilag (1979): Fruit rot of pepper caused
by Fusarium spp. Philippine Phytopathology (1977): 13
(1/2): 14-23.

Mitchell, F.G. J.P. Gentry, R. Guillow and M.H. Gordts (1964):
Tight fill peach packing- laboratory and transit tests.
Calif Agric. 18 (11): 13-14.

Mitchell, F.G., R. Guillow and R.A. Persons (1972): Commercial
cooling of fruits and vegetables. Calif Agric Exp. Sta.
Ext. Serv. Man. 43-44 pp.

Mitchell, F.G., W.C. Mieke, F.P. Guerreiro and G. Mayor (1967):
Packing sweet cherries to reduce transit injury. Calif
Agric. 21 (8): 6-7.

Morris, L.L. and A.A. Kador (1977): Commodity requirements and
recommendations for transport and storage selected
vegetables. Mich. State Univ. Hortic. Rept. 28, pp
266-76.

Morris, L.L., L.L. Claypool and D.P. Murr (1971): Modified
atmospheres - an Indexed Reference List Through 1969,
with emphasis on Horticultural Commodities, Berkely:
Univ Calif Press 115 pp.

- Mullen, J.M. and D.F. Bateman (1976): Polysaccharide degrading enzymes produced by Fusarium roseum "Avenaceum" in culture and during pathogenesis.
- Mukerjee, P.K. (1961): Cold storage of mangoes. Gardening 3 : 53-54.
- Naik, R. (1979): A fruit rot of apple caused by Monochaetia unii Science & Culture (1977): 43 (12) 551-552 Univ Illinois Urbana U.S.A.
- Naqvi, S.H.Z. & H.M. Good (1957): Studies of the aging of conidia of Monilinia fructicola (Wint) Honey. I Germination rates and longevity. Can. J. Botany 35: 635-645.
- Naranjo, K. and S.M. Reddy (1980): Studies on fusarial rot of lemons. Indian Phytopathology (1978, publ. 1979): 31 (3): 310-314. Kakatiya Univ. Warangal, A.P. India.
- Olsen, E.L., G.O. Patchen and H.A. Schomer (1960): Cooling rates of apples packed in fiber board boxes as influenced by vents, perforated trays, and stacking pattern. Wash. State. Hortic. Assoc. Proc. 36: 214-20.
- Onesiroso, P.T. and T. Fatunla (1977): Fungal fruit rots of tomatoes in Southern Nigeria. J. Hortic. Sci. 51 (4): 473-479.
- Ooka, J.J. (1981): Guava fruit rot caused by Rhizopus stolonifer in Hawaii. Plant Disease (1980): 64 (4) 412-413. Univ. Hawaii, Kaula Branch St, Kapaa, Hawaii, U.S.A.
- Pandey, R.S., D.N. Shukla; D.V.S. Khatri and S.N. Bhargava (1980):

- A new fruit rot of Phyllanthus emblica. Indian Phytopath. **33** (3) 491.
- Pandey, R.S., D.V.S. Khati, D.N. Shukla and S.N. Bhargava (1980): A new Phoma rot of bael fruit. Indian Phytopath. **33** (2): 319.
- Pantastico, E.B. (1975): Post harvest Physiology, Handling and Utilization of Tropical and Sub-Tropical Fruits and Vegetables. Westport, Conn AVI Publ 360 pp.
- Panwar, N.S., J.N. Chand, Harnam Singh and C.S. Paracer (1972): Phomopsis fruit rot of brinjal (Solanum melongena L.) in Punjab. I. Viability of the fungus and role of seeds in disease development. J. Res. Punjab Agric. Univ. **7** 641-643.
- Parpia, H.A.B. (1976): Post harvest losses impact of their prevention on food supplies, nutrition and development. In Symp on Nutrition and Agricultural and Economics Development in the Tropics. Guatemala, 1974 ed N.S. Srinashaw, M. Behar, pp 195-206. New York: Plenum 500 pp.
- Patchen, G.O. (1969): Effects of vent holes on strength of fiberboard boxes and fruit cooling rate. US De. Agric ARS 52-34 12 pp.
- Pearson, R.C. and D.H. Hall (1974): Ripe fruit rot of tomato caused by Pythium ultimum and Pythium achanidomaticum Plant Dis Rep. **57** (12): 1066-1069.
- Pendergrass, A., F.M.R. Isenberg (1974): The effect of

relative humidity on the quality of cabbage. Hort Science 9: 226-27.

Perkins, F.A. (1959): Bagged apple damage studied - Prod. Marks 2 (4): 27.

Philip, S. (1981): Aspergillus rot of pomegrate fruits. India Phytopathology (1979, publ. 1980); 32 (2) 332. Coll. Agric. Vellayani, Kerala India.

Phillips, D.J. and G.M. Harris (1980): Post harvest blown rot of peaches and inoculum density of Monilia fructicola (Wint). Agricultural Research Results, Science and Education Administration Western Series (1979); 9 12 pp.

Powell, G.H. (1960): The handling of fruit for transportation U.S.D.A. Irr. Book 1905: 329-336.

Prakash, O.M., B. Mishra and A.P. Mishra (1976): Greenaria fuliginosa Scribner and Viala causing bitter rot of grapes in India. Indian Phytopathology 27 (4): 603-606.

Prasad, S.S. and S.K. Sinha (1979): Post infection changes in ascorbic acid content of litchi fruits caused by certain pathogenic fungi. Proceedings of the National Academy of Sciences India B (1975); 45(1) 43-44.

Popkova, K.V. and L.V. Kovaleva (1974): Ophiobolus porphyreus (Tode) Sacc, the anamycetous stage of the fungus causing potato rot. Mikol. Fitopatol. 7 (3): 233-237.

- Popov, F.A. (1979): Biological features of the pathogen of Phoma rot of potato (Phoma solanicola Prill and Dell).
From Referativnyi Zhurnal Biologii (1978); 9V 307.
- Quimio, T.H. and A.J. Quimo (1977): A new fruit rot disease of papaya in the Philippines: PHILIPP AGRIC; 58 (7/8): 330-331 1974/1975.
- Quimio, T.H. & L.E. Abilay (1981): Corynespora disease of papaya in the philippines. Philippine Phytopathology; (1979); 15 (2): 158-161. Univ. Philippines, Los Banos, College Laguna, Philippines.
- Rai, R.P. (1980): Fusarium equiseti (Corda) Sacc causing dry rot of potato tubers - a new report Current Science (1979); 48 (23): 1043-1045 Cent. Potato, Res. Inst. Simla, India.
- Rajagopalan, B. and K.I. Wilson (1972): Diplodia natalensis causing dry rot of guava fruits in S. India. Plant Dis Rep 56 (4): 323-324.
- Rajak, R.C. and S.P. Gautam (1981): Two new fruit rot disease of Citrus. Current Science (1980); 42 (15): 597-598. Univ. Jabalpur, M.P. India.
- Raju, K.S. and R.B. Raj (1981): Soft rot of onion in storage. Madras Agricultural Journal. (1980) 67 (3): 194-195. Agric. Res. Sta. Lam. A.P. India.
- Rath, G.C. and G.N. Mohanty (1980): Control of Macrophoma rot of stored garlic. Indian Phytopathology (1978, publ. 1979); 31 (2) 256-257 Orissa Univ. Agric. Technol.

Bhubaneswar India.

- Rao, S.S. and V. Thirupathaiah (1980): Penicillium lanthionellum causing soft rot of snake gourd (Trichosanthes amarus). Indian Phytopathology (1978, publ. 1979); 21 (4): 532. Kakatiya Univ. Warangal, India.
- Rao, V.G. and V. Subramaniam (1978): A new post harvest disease of cape-gooseberry Journal of the University of Bombay Sciences (1976); 45 (72): 58-61. Maharashtra Ass. Cultivation Sci, Poona, India.
- Redit, W.H. (1969): Protection of rail shipments of fruits and vegetables. US Dep Agric Handb. 193. 98 pp.
- Rondomanski, W. (1978): White rot of Onion (Sclerotium cepivorum Berk) in Poland. Polskie Biuletyn Instytutu Ochrony Roslin (1974) 37, 187-190.
- Roy, A.N.; G. Sharma and R.B. Sharma (1981): Penicillium dry fruit rot of ashgourd (Penicillium citrinum) on Benincasa hispida. Current Science, (1980); 49 (17): 670. Agra Coll, India.
- Roy, A.N. and R.B. Sharma (1980): / fruit rot of Pyrus communis caused by Trichoderma viride. Current Science, (1979); 48 (11): 498-499. Agra Coll. India.
- Roy, A.N. R.B. Sharma and K.C. Gupta (1978): Occurrence of 3 new rot diseases of stored garlic (Allium sativum L.) Current Science, (1977) 46 (20) 716-717 Agra Coll. U.P. India.

- Rygg, G.L., A.W. Wells, S.M. Norman and E.P. Atrops (1964):
Biphenyl control of citrus spoilage - influence of time,
temperature and carton venting. US Dep. Agric. Mark. Res.
Rept. 646. 22 pp.
- Saxena, A.K., and E.K. Jain (1981): A new fruit rot disease of
papaya (Trichothecium roseum on papaw). Current Science,
(1981) 50 (2): 96 (Eng) Jivaji Univ. Gwalior, India.
- Scheffrath, J. (1978): Damage by Chalaropsis thielavicioides
Peyr. on stored carrots in the Frankfurt district.
Nachrichtenblatt für den Pflanzenschutz in der DDR,
(1978), 32 (4) 83.
- Seenappa, M.; L.W. Stobbs and A.C. Kempton (1981): Asporangium
colonization of Indian red pepper during storage. Phytopathology, (1980), 70 (3): 218-222. Univ. Waterloo, Ont.
Canada.
- Sethi, D.R. (1963): Cold storage of seed potatoes. Indian Mag.
9 : 471-473.
- Sharma, N. and A.M. Khan (1978): Fruit rot of Averrhoa caram-
bola L. Cliodsporium herbarum Geoplos. (1978); 3 (1): 48,
A.M.U. Aligarh, India.
- Sharma, N. and A.M. Khan (1979): Studies on fruit rot of cape
gooseberry. Acta Botanica India, (1978); 6 (2): 211-213
A.M.U., Aligarh, India.

(Amorphophallus semipalmatus Blume) caused by Sclerotium rolfsii Sacc. Science and Culture (1977); 43 (11): 508.
Agric. Res. Sta. Borkhara Farm, Kota, India.

Upstone, M.E. (1979): Phytophthora syringae fruit rot of apples; Plant Pathology (1978); 27 (1): 24-30.ADAS, Wye,
Ashford, Kent, U.K.

Varkhrushova, T.E. (1973): Causative agent of carrot black rot
Mikol Fitopatol. 61: 48-50.

Verma, A.S.; A. Cheeran and C.K. Peethambaram (1981): Fruit rot
of bitter gourd. (Glomerella singulata on Momordica
charantia). Agricultural Research Journal of Kerala,
(1980) 16 (1): 135-136. Coll. Hort. Vellanikkara, Trichur,
Kerala India.

Verma, R.A.B.; J.S. Prasad and N.K. Jha (1977): Investigation on
the Alternaria rot of akhrot (Juglans regia-L) Sydney
Ann Mysol; 26 (1-6): 158-161.

Vyas, K.M.; S.C. Chaurasia and N. Saxena (1980): A new fruit
rot disease of brinjal caused by Rhizopus nodosus. Indian
Phytopathology; (1978, publ. 1979) 31 (3): 363. Univ.
Saugar, Sagar M.P. India.

Wadia, K.D.R. and C. Manoharachary (1979): The new post-harvest
disease of Carissa carandas L. Fruits. Geobion, (1979) 6
(3): 126-127. Sci- Coll. Osmania Univ. Hyderabad India.

Wadia, K.D.R. and C. Manoharachary (1981): Three post-harvest
diseases of Zizyphus mauritiana fruits from India. Plant

disease (1980); 64 (3): 323-325. Sci. Coll. Sairabad
Osmania Univ Hyderabad A.P. India.

Wasfy, E.H. H.M. Elarosi; S.H. Michail and M.K. Tawood (1978):
Pathogens of cucumber fruit rots in Egypt. Egyptian
Journal of Phytopathology (1975) 7: 67-70. Alexandria
Univ. Egypt.

Wells, J.M. (1971): Post-harvest hot water and fungicide
treatments for reduction of decay of California peaches,
plums and nectarines. US Dept. Agric. Mark. Res. Rept.
908, 12pp.

Weresub, L.K. and W.L. Illman (1980): Corticium centrifugum
reisolated from filchey rot of stores apples. Canadian
Journal of Botany (1980); 58 (2): 137-146. Biosystoma-
tica Res Inst Agric Canada, Ottawa, Ont, Canada.

Whitman, T.M. (1957): Freezing points of fruits, vegetables and
florist stocks. US Dep. Agric. Mark. Res. Rept. 196
32 pp. Reprinted 1977.

Williamson, D. and R.N. Tandon (1965): Naturwissenschaften 7: 166

Wilson, K.I. and C. Balagopal (1973): Dry rot of ginger rhizomes
caused by Diplodia natalensis. Indian Phytopathol. 24 (2):
385-386.

Wu, W.S. (1978): Black mold rot of onion in Taiwan. Plant Pro-
tection Bulletin Taiwan. (1977); 19 (3): 202-205. Natn.
Taiwan Univ Taipei, Taiwan.